

# SEARCH REQUEST FORM

Scientific and Technical Information Center

Access DB# 94863  
RECEIVED

JUN 18 2003

Requester's Full Name: RGITOMEN Examiner #: 69630 (STIC) Date: 6/18/03  
Art Unit: 1651 Phone Number 301-8-0732 Serial Number: 09/977,667  
Mail Box and Bldg/Room Location: 11301 Results Format Preferred (circle): PAPER DISK E-MAIL  
11D11

If more than one search is submitted, please prioritize searches in order of need.

\*\*\*\*\*  
Please provide a detailed statement of the search topic, and describe as specifically as possible the subject matter to be searched. Include the elected species or structures; keywords, synonyms, acronyms, and registry numbers, and combine with the concept or utility of the invention. Define any terms that may have a special meaning. Give examples or relevant citations, authors, etc, if known. Please attach a copy of the cover sheet, pertinent claims, and abstract.

Title of Invention: \_\_\_\_\_

Inventors (please provide full names): \_\_\_\_\_

Earliest Priority Filing Date: \_\_\_\_\_

\*For Sequence Searches Only\* Please include all pertinent information (parent, child, divisional, or issued patent numbers) along with the appropriate serial number.

JAN

Jan Delaval  
Reference Librarian  
Biotechnology & Chemical Library  
CM1 1E07 - 703-308-4498  
jan.delaval@uspto.gov

## STAFF USE ONLY

	Type of Search	Vendors and cost where applicable
Searcher: <u>Jan</u>	NA Sequence (#) _____	STN <input checked="" type="checkbox"/>
Searcher Phone #: <u>4498</u>	AA Sequence (#) _____	Dialog <input checked="" type="checkbox"/>
Searcher Location: _____	Structure (#) _____	Questel/Orbit _____
Date Searcher Picked Up: <u>6/30/03</u>	Bibliographic <input checked="" type="checkbox"/>	Dr.Link _____
Date Completed: <u>6/30/03</u>	Litigation _____	Lexis/Nexis _____
Searcher Prep & Review Time: _____	Fulltext _____	Sequence Systems _____
Clerical Prep Time: <u>15</u>	Patent Family _____	WWW/Internet _____
Online Time: <u>5:50</u>	Other _____	Other (specify) _____

PTO-1590 (8-01)

=> fil hcaplus

FILE 'HCAPLUS' ENTERED AT 09:54:14 ON 30 JUN 2003

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

Jan Delaval  
Reference Librarian  
Biotechnology & Chemical Library  
CM1 1E07 - 703-308-4498  
jan.delaval@uspto.gov

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 30 Jun 2003 VOL 139 ISS 1  
FILE LAST UPDATED: 29 Jun 2003 (20030629/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d all hitstr tot

L69 ANSWER 1 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 2003:320162 HCAPLUS

DN 138:299800

TI A system for detection of **urease** in a human **gastric sample** for diagnosis of **gastrointestinal** bacterial infection

IN **McMichael, Donald J.; Peterson, Kristy; Marshall, Barry J.; Mendis, Aruni H. W.; Chairman, Simon**

PA **Kimberly-Clark Worldwide, Inc., USA**

SO PCT Int. Appl., 34 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM G01N033-48

CC 7-1 (Enzymes)

Section cross-reference(s): 14

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003034061	A2	20030424	WO 2002-US29814	20020918
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	US 2003077680	A1	20030424	US 2001-977555	20011015
	US 2003077684	A1	20030424	US 2001-977874	20011015
	US 2003082664	A1	20030501	US 2001-977556	20011015
	US 2003082661	A1	20030501	US 2001-977667	20011015
PRAI	US 2001-977555	A	20011015		
	US 2001-977556	A	20011015		

US 2001-977667 A 20011015


US 2001-977874 A 20011015

- AB A system and method for detecting bacterial infections in the human **gastrointestinal** tract is disclosed. In one embodiment, the system includes a first compn. sepd. from a second compn. The first compn. contains **urea** in **powd.** form. The second compn., on the other hand, contains an **indicator**. A **biopsy** of a **gastric sample** is first contacted with the first compn. and then placed in the second compn. The second compn. **indicates** the presence of an enzyme that, in turn, **indicates** the presence of bacteria. In an alternative embodiment of the present invention, a **biopsy** of a **gastric sample** is contacted with a single compn. The compn. contains **urea** in a **powd.** form combined with a dry **indicator**. Besides **urea** and a dry **indicator**, the compn. can also contain an anticaking agent. The system of the present invention can include a container for holding the compns. A specimen handling tool can be included in the container for handling a **biopsy sample**.
- ST **urease biopsy detection gastrointestinal**  
bacterial infection diagnosis human; **urea** anticaking agent  
**indicator urease detn gastrointestinal**  
infection diagnosis
- IT Titration  
(acid-base, **pH** adjuster; system for detection of  
**urease** in human **gastric sample** for  
diagnosis of **gastrointestinal** bacterial infection)
- IT Analytical apparatus  
(biochem.; system for detection of **urease** in human  
**gastric sample** for diagnosis of  
**gastrointestinal** bacterial infection)
- IT Stomach  
(**biopsy**; system for detection of **urease** in human  
**gastric sample** for diagnosis of  
**gastrointestinal** bacterial infection)
- IT Digestive tract, disease  
(infection; system for detection of **urease** in human  
**gastric sample** for diagnosis of  
**gastrointestinal** bacterial infection)
- IT Diagnosis  
(mol.; system for detection of **urease** in human  
**gastric sample** for diagnosis of  
**gastrointestinal** bacterial infection)
- IT Particle size  
(of **urea powder**; system for detection of  
**urease** in human **gastric sample** for  
diagnosis of **gastrointestinal** bacterial infection)
- IT **Acid-base indicators**  
Agglomeration preventers  
Antibacterial agents  
**Colorimetric indicators**  
Containers  
Digestive tract  
Digestive tract, disease  
Films  
**Gels**  
Human  
**Indicators**  
**Powders**  
**Sample preparation**  
Test kits  
(system for detection of **urease** in human **gastric**  
**sample** for diagnosis of **gastrointestinal** bacterial

- infection)
- IT 1344-00-9, Sodium aluminosilicate 7631-86-9, Silica, biological studies  
 RL: ARU (Analytical role, unclassified); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (anticaking agent; system for detection of **urease** in human **gastric sample** for diagnosis of **gastrointestinal** bacterial infection)
- IT 9002-18-0, Agar  
 RL: ARU (Analytical role, unclassified); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (gel; system for detection of **urease** in human **gastric sample** for diagnosis of **gastrointestinal** bacterial infection)
- IT 57-13-6, Urea, biological studies  
 RL: ARG (Analytical reagent use); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (powder; system for detection of **urease** in human **gastric sample** for diagnosis of **gastrointestinal** bacterial infection)
- IT 7664-41-7, Ammonia, biological studies  
 RL: ANT (Analyte); ARG (Analytical reagent use); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (system for detection of **urease** in human **gastric sample** for diagnosis of **gastrointestinal** bacterial infection)
- IT 9002-13-5, Urease  
 RL: ANT (Analyte); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (system for detection of **urease** in human **gastric sample** for diagnosis of **gastrointestinal** bacterial infection)
- IT 143-74-8, Phenol red  
 RL: ARG (Analytical reagent use); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (system for detection of **urease** in human **gastric sample** for diagnosis of **gastrointestinal** bacterial infection)
- IT 9002-18-0, Agar  
 RL: ARU (Analytical role, unclassified); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (gel; system for detection of **urease** in human **gastric sample** for diagnosis of **gastrointestinal** bacterial infection)
- RN 9002-18-0 HCAPLUS  
 CN Agar (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

- IT 57-13-6, Urea, biological studies  
 RL: ARG (Analytical reagent use); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (powder; system for detection of **urease** in human **gastric sample** for diagnosis of **gastrointestinal** bacterial infection)
- RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)



- IT 7664-41-7, Ammonia, biological studies

RL: ANT (Analyte); ARG (Analytical reagent use); DGN (Diagnostic use);  
ANST (Analytical study); BIOL (Biological study); USES (Uses)  
(system for detection of **urease** in human **gastric**  
**sample** for diagnosis of **gastrointestinal** bacterial  
infection)

RN 7664-41-7 HCAPLUS  
CN Ammonia (8CI, 9CI) (CA INDEX NAME)

NH<sub>3</sub>


IT 9002-13-5, Urease  
RL: ANT (Analyte); DGN (Diagnostic use); ANST  
(Analytical study); BIOL (Biological study); USES (Uses)  
(system for detection of **urease** in human **gastric**  
**sample** for diagnosis of **gastrointestinal** bacterial  
infection)

RN 9002-13-5 HCAPLUS  
CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

IT 143-74-8, Phenol red  
RL: ARG (Analytical reagent use); DGN (Diagnostic use); ANST (Analytical  
study); BIOL (Biological study); USES (Uses)  
(system for detection of **urease** in human **gastric**  
**sample** for diagnosis of **gastrointestinal** bacterial  
infection)

RN 143-74-8 HCAPLUS  
CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis- (9CI) (CA  
INDEX NAME)



L69 ANSWER 2 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 2003:76201 HCAPLUS

DN 138:300142

TI Method for determining Helicobacter pylori-associated **intra**gastral  
**urease** activity from **biopsies**

IN Nizhevich, A. A.; Sataev, V. U.; Khasanov, R. Sh.; Mel'nikova, Z. M.;  
Loginovskaya, V. V.; Akhmetshin, R. Z.

PA Bashkirskii Gosudarstvennyi Meditsinskii Universitet, Russia

SO Russ., No pp. given

CODEN: RUXXE7

DT Patent

LA Russian

IC ICM G01N033-48

ICS G01N033-49

CC 9-4 (Biochemical Methods)

## Section cross-reference(s): 14

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	RU 2189591	C1	20020920	RU 2001-102196	20010124
PRAI	RU 2001-102196		20010124		

AB The inventive method deals with detecting the degree of bacterial seeding vol. of **gastric** mucosa at **Helicobacter gastritis**, **gastroduodenitis** and ulcerous disease. One should take a **biopsy** fragment of **gastric** mucosa and put it into com. soln. Incubation mixt. is subjected for exposure, **PEC-colorimetry** is conducted at 540 nm wave length to compare optic d. with incubation time of **biopsy** fragment and the wt. of **biopsy** fragment (units of optic d./mg **biopsy** material/min). At **urease** activity values ranged 11.5-4 U one should detect a low degree of bacterial seeding vol. of **gastric** mucosa, at its value within 5-10 U a moderate degree is detected and in case its value ranges 11-19 U a high degree of **H.pylori** seeding vol. is concluded on. The method is of high specificity, enables to conduct a semi-quant. anal. of bacterial seeding vol. of **gastric** mucosa.

ST **Helicobacter** assocd **intra**gastral **urease** activity **colorimetry biopsy**

IT **Intestine**, disease  
(duodenum, ulcer; method for detg. **Helicobacter pylori**-assocd. **intra**gastral **urease** activity from **biopsies**)

IT Stomach, disease  
(**gastritis**; method for detg. **Helicobacter pylori**-assocd. **intra**gastral **urease** activity from **biopsies**)

IT **Colorimetry**  
**Helicobacter pylori**  
(method for detg. **Helicobacter pylori**-assocd. **intra**gastral **urease** activity from **biopsies**)

IT Stomach  
(mucosa; method for detg. **Helicobacter pylori**-assocd. **intra**gastral **urease** activity from **biopsies**)

IT 9002-13-5, **Urease**  
RL: **ANT (Analyte)**; **DGN (Diagnostic use)**; **ANST (Analytical study)**; **BIOL (Biological study)**; **USES (Uses)**  
(method for detg. **Helicobacter pylori**-assocd. **intra**gastral **urease** activity from **biopsies**)

IT 57-13-6, **Urea**, biological studies  
RL: **ARG (Analytical reagent use)**; **DGN (Diagnostic use)**; **ANST (Analytical study)**; **BIOL (Biological study)**; **USES (Uses)**  
(method for detg. **Helicobacter pylori**-assocd. **intra**gastral **urease** activity from **biopsies**)


IT 9002-13-5, **Urease**  
RL: **ANT (Analyte)**; **DGN (Diagnostic use)**; **ANST (Analytical study)**; **BIOL (Biological study)**; **USES (Uses)**  
(method for detg. **Helicobacter pylori**-assocd. **intra**gastral **urease** activity from **biopsies**)

RN 9002-13-5 HCAPLUS  
CN **Urease** (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

IT 57-13-6, **Urea**, biological studies  
RL: **ARG (Analytical reagent use)**; **DGN (Diagnostic use)**; **ANST (Analytical study)**; **BIOL (Biological study)**; **USES (Uses)**  
(method for detg. **Helicobacter pylori**-assocd. **intra**gastral **urease** activity from **biopsies**)

RN 57-13-6 HCAPLUS  
CN **Urea** (8CI, 9CI) (CA INDEX NAME)



L69 ANSWER 3 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 2003:24772 HCAPLUS

DN 138:217792

TI Method and test kit of diagnosing helicobacteriosis from estimation of **urease** activity of biological material

IN Dmitrienko, M. A.; Kornienko, E. A.; Mileiko, V. E.

PA Russia

SO Russ., No pp. given

CODEN: RUXXE7

DT Patent

LA Russian

IC ICM C12Q001-04

ICS C12Q001-00

CC 9-1 (Biochemical Methods)

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	RU 2184781	C2	20020710	RU 1997-117123	19970930
PRAI	RU 1997-117123		19970930		

AB The invention relates to investigations involving control of **urease** activity of tissue **samplings** and body fluids in order to est. their bacterial loading, in particular by *Helicobacter pylori*. **Urease** activity is detd. from the value or time of emergence of **indication** effect of **color** reaction proceeding on solid sorbent as the result of interaction of acid-base **indicator** and products of **urea-to-ammonia** hydrolysis caused by endogenous **urease** of microorganisms. **Ammonia** is fixed by solid capillary or grainy sorbent. Method is realized in device allowing performing express anal. of tissue **samplings**, body fluids, and aerosols. **Urea** and acid-base **indicator** are deposited on solid hygroscopic fibrous or grainy microcapillary sorbent in the form of homogeneous fine-crystal dispersion. Tissue **samplings** can also be used for other investigations.

ST *Helicobacter helicobacteriosis urease* detn test kit

IT Bacteria (Eubacteria)

(helio-, infections; method and test kit of diagnosing helicobacteriosis from estn. of **urease** activity of biol. material)

IT Absorbents

**Acid-base indicators**

Body fluid

**Colorimetry**

*Helicobacter pylori*

**Sampling**

Test kits

(method and test kit of diagnosing helicobacteriosis from estn. of **urease** activity of biol. material)

IT 9002-13-5, **Urease**

RL: ANT (Analyte); DGN (Diagnostic use); ANST

(Analytical study); BIOL (Biological study); USES (Uses)

(method and test kit of diagnosing helicobacteriosis from estn. of **urease** activity of biol. material)

IT 57-13-6, **Urea**, biological studies


RL: ARG (Analytical reagent use); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(method and test kit of diagnosing helicobacteriosis from estn. of

urease activity of biol. material)  
 IT 9002-13-5, Urease  
 RL: ANT (Analyte); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (method and test kit of diagnosing helicobacteriosis from estn. of urease activity of biol. material)  
 RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

IT 57-13-6, Urea, biological studies  
 RL: ARG (Analytical reagent use); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (method and test kit of diagnosing helicobacteriosis from estn. of urease activity of biol. material)  
 RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)



L69 ANSWER 4 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 2002:872769 HCAPLUS

DN 137:334615

TI Method and device for urease determination in roasted soybean

IN Destri, Maurizio; Maradini, Claudio; Marocchi, Daniela

PA Raggio Di Sole Mangimi S.P.A., Italy

SO Ital., 11 pp.

CODEN: ITXXBY

DT Patent

LA Italian

IC ICM A23K

ICS G01N

CC 7-1 (Enzymes)

Section cross-reference(s): 11, 17

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	IT 1304527	B1	20010319	IT 1998-PR30	19980511
PRAI	IT 1998-PR30		19980511		
AB	A method and device for urease detn. in exts. of roasted soybeans is disclosed. The ext. is combined with a soln. contg. urea and the amt. of ammonia released is measured with pH paper. A tube for the substrate soln. and soy ext. as well as a stopper contg. pH paper which is used in the urease detn. is also disclosed.				
ST	soybean soy flour urease detn app ammonia pH paper				
IT	Apparatus				
	(method and device for urease detn. in roasted soybean)				
IT	Acid-base indicators				
	(pH paper; method and device for urease detn. in roasted soybean)				
IT	Soybean (Glycine max)				
	(roasted; method and device for urease detn. in roasted soybean)				
IT	7664-41-7, Ammonia, biological studies				
	RL: BSU (Biological study, unclassified); BIOL (Biological study)				
	(detection with pH paper of; method and device for				




urease detn. in roasted soybean)  
 IT 9002-13-5, Urease  
 RL: ANT (Analyte); ANST (Analytical study)  
 (method and device for urease detn. in roasted soybean)  
 IT 57-13-6, Urea, biological studies  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (substrate; method and device for urease detn. in roasted  
 soybean)  
 IT 7664-41-7, Ammonia, biological studies  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (detection with pH paper of; method and device for  
 urease detn. in roasted soybean)  
 RN 7664-41-7 HCAPLUS  
 CN Ammonia (8CI, 9CI) (CA INDEX NAME)

NH<sub>3</sub>

IT 9002-13-5, Urease  
 RL: ANT (Analyte); ANST (Analytical study)  
 (method and device for urease detn. in roasted soybean)  
 RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

IT 57-13-6, Urea, biological studies  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (substrate; method and device for urease detn. in roasted  
 soybean)  
 RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)



L69 ANSWER 5 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
 AN 2002:125559 HCAPLUS  
 DN 136:130760  
 TI Detection method of Helicobacter pylori using rapid urease  
 detection kit  
 IN Lee, Jong Wook; Kim, Beom Su; Bae, Su Hwan; Lee, Gyeong Won; Jeong, Yun  
 Seob  
 PA S. Korea  
 SO Repub. Korean Kongkae Taeho Kongbo, No pp. given  
 CODEN: KRXXA7  
 DT Patent  
 LA Korean  
 IC ICM C12Q001-04  
 CC 7-1 (Enzymes)  
 Section cross-reference(s): 10, 14

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI KR 2000033013	A	20000615	KR 1998-49668	19981119
PRAI KR 1998-49668		19981119		
AB	PURPOSE: Detection kit of Helicobacter pylori using rapid urease test is provided which is specific and sensitive to detect urease of Helicobacter pylori in the tissue of stomach from human or animals.			

CONSTITUTION: **Ammonia** prodn. from the tissue of stomach which is suspended in HCl-KCl **buffer** not being added **urea** is high enough to detect. Addn. of small amt. of **urea** into the **buffer** saves the time of test and increases the sensitivity of the test. A test kit comprises acid **buffer** (pH 2.0-5.0) and **indicator**. The acid **buffer** consists of HCl and KCl. Congo red is used as an **indicator** and the **color** change from blue to red is pos. sign. The optimal concn. of **indicator** ranges from 50mg/mL to 2g/mL. The amt. of **urea** added into **buffer** is 50-500mg/mL. The test is performed by only suspension of test tissue in the kit or shaking of the tube.

ST

IT

Animal tissue  
**Buffers**  
**Colorimetry**  
 Concentration (condition)  
 Helicobacter pylori  
 Human

#### **Indicators**

Mixing  
 Stomach  
 Suspensions  
 Test kits  
 Time

#### **pH**

(detection method of Helicobacter pylori using rapid **urease** detection kit)

IT

Acids, biological studies

RL: ARU (Analytical role, unclassified); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(detection method of Helicobacter pylori using rapid **urease** detection kit)

IT

14798-03-9, Ammonium, biological studies

RL: ANT (Analyte); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study)

(detection method of Helicobacter pylori using rapid **urease** detection kit)

IT

9002-13-5, Urease

RL: ANT (Analyte); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(detection method of Helicobacter pylori using rapid **urease** detection kit)

IT

57-13-6, Urea, biological studies 573-58-0, Congo red

RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(detection method of Helicobacter pylori using rapid **urease** detection kit)

IT

7447-40-7, Potassium chloride (KCl), biological studies 7647-01-0, Hydrogen chloride, biological studies

RL: ARU (Analytical role, unclassified); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(detection method of Helicobacter pylori using rapid **urease** detection kit)

IT

9002-13-5, Urease

RL: ANT (Analyte); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(detection method of Helicobacter pylori using rapid **urease** detection kit)


RN

9002-13-5 HCAPLUS

CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

IT 57-13-6, Urea, biological studies  
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST  
 (Analytical study); BIOL (Biological study); USES (Uses)  
 (detection method of Helicobacter pylori using rapid urease  
 detection kit)  
 RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)



L69 ANSWER 6 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 2001:851420 HCAPLUS

DN 135:355035

TI Novel method for the isolation of Helicobacter pylori from highly  
 contaminated specimens

IN Song, Qunsheng; Zirnstein, Gerald W.; Gold, Benjamin D.

PA United States Dept. of Health and Human Services, USA

SO PCT Int. Appl., 41 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM C12Q001-00

CC 9-16 (Biochemical Methods)

Section cross-reference(s): 10

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001088183	A2	20011122	WO 2001-US40756	20010516
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 2001059869	A5	20011126	AU 2001-59869	20010516
PRAI US 2000-205320P	P	20000518		
WO 2001-US40756	W	20010516		

AB Methods and kits are disclosed for isolating urease-pos.  
 bacteria by exposing a sample for 1 to 60 min to a media contg.  
 urea along with simultaneous or subsequent exposure to pH  
 below 3.0. In one embodiment, the bacteria is H. pylori and the acidic  
 conditions are provided by addn. of HCl. These methods and kits are esp.  
 useful for isolating or detecting H. pylori in samples, such as  
 saliva samples, contaminated by other microorganisms.

ST isolation Helicobacter pylori contaminated specimen

IT Medical goods

(Endoscopy equipment; novel method for isolation of Helicobacter pylori  
 from highly contaminated specimens)

IT Stomach

(biopsy; novel method for isolation of Helicobacter pylori  
 from highly contaminated specimens)

IT Filters

(hydrophobic; novel method for isolation of Helicobacter pylori from

highly contaminated specimens)

IT Bacteria (Eubacteria)  
 Blood  
 Cell  
 Concentration (condition)  
 Culture media  
 Environment  
 Feces  
 Filters  
 Filtration  
 Growth, microbial  
 Helicobacter pylori  
 Microorganism  
 Proteus (bacterium)  
 Rabbit  
 Saliva  
**Samples**  
 Test kits  
 Time  
**pH**  
 (novel method for isolation of Helicobacter pylori from highly contaminated specimens)

IT Acids, biological studies  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
 (Uses)  
 (novel method for isolation of Helicobacter pylori from highly contaminated specimens)

IT Tooth  
 (plaque; novel method for isolation of Helicobacter pylori from highly contaminated specimens)

IT **7664-41-7, Ammonia**, analysis  
 RL: ANT (Analyte); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study)  
 (novel method for isolation of Helicobacter pylori from highly contaminated specimens)

IT **9002-13-5, Urease**  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (novel method for isolation of Helicobacter pylori from highly contaminated specimens)

IT **57-13-6, Urea**, biological studies 7647-01-0, Hydrogen chloride, biological studies **9002-18-0, Agar**  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
 (Uses)  
 (novel method for isolation of Helicobacter pylori from highly contaminated specimens)

IT **7664-41-7, Ammonia**, analysis  
 RL: ANT (Analyte); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study)  
 (novel method for isolation of Helicobacter pylori from highly contaminated specimens)

RN 7664-41-7 HCAPLUS  
 CN Ammonia (8CI, 9CI) (CA INDEX NAME)

NH3

IT **9002-13-5, Urease**  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (novel method for isolation of Helicobacter pylori from highly contaminated specimens)

RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

IT 57-13-6, Urea, biological studies 9002-18-0,


Agar

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)

(novel method for isolation of Helicobacter pylori from highly  
contaminated specimens)

RN 57-13-6 HCAPLUS

CN Urea (8CI, 9CI) (CA INDEX NAME)



RN 9002-18-0 HCAPLUS

CN Agar (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 7 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 2001:582293 HCAPLUS

DN 135:133929

TI Detection of H. pylori in the stomach

IN Marshall, Barry

PA Australia

SO U.S. Pat. Appl. Publ., 6 pp., Cont.-in-part of U.S. 6,228,605.

CODEN: USXXCO

DT Patent

LA English

IC ICM C12Q001-04

NCL 435034000

CC 7-1 (Enzymes)

Section cross-reference(s): 10, 14

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2001012623	A1	20010809	US 2001-824870	20010403
PRAI	US 1995-489816	B1	19950613		
	US 1997-832332	A2	19970326		

AB A method for the in vivo detection of **urease**-producing Helicobacter in the upper stomach is disclosed. The dense carrier is divided into two sep. groups which are combined with sep. reagent **indicators**, one of which also contains **urea**. The carriers are food sol. products, preferably sugar beads having a diam. of approx. 0.2 to 3.0 mm. The treated carriers and **urea** are encapsulated in a sol. capsule which is administered to a patient. The d. of the carriers cause the capsule to migrate to the **gastric** mucosa, where the capsule, but not the reagents, is dissolved, placing the reagents and **urea** in direct contact with the **gastric** mucosa. The **urea** reacts with any **urease** present in the stomach by creating **ammonia**, which increases the **pH** in the immediate vicinity of the **urea** contg. carrier and **indicator** beads. The two reagents react differently, through **color** change, to the increase in **pH**, which is viewed through use of an endoscope. A preferred first reagent is bromothymol blue (dibromothymolsulfonphthalein), which changes yellow in the presence of **urease**, and a preferred second reagent is **phenol red** (phenolsulfonphthalein), which turns red in the presence of **urease**.

ST detection Helicobacter pylori stomach

IT Capsules  
(Sol.; detection of H. pylori in stomach)

IT Spheres  
(beads; detection of H. pylori in stomach)

IT Carriers  
Colorimetry  
Encapsulation  
Endoscopes  
Food  
Gastric juice  
Helicobacter pylori  
Stomach  
Stomach content  
pH  
(detection of H. pylori in stomach)

IT Reagents  
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)  
(detection of H. pylori in stomach)

IT Carbohydrates, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(detection of H. pylori in stomach)

IT Stomach  
(mucosa; detection of H. pylori in stomach)

IT 9002-13-5, Urease  
RL: ANT (Analyte); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
(detection of H. pylori in stomach)

IT 57-13-6, Urea, uses 76-59-5, Bromothymol blue  
143-74-8, Phenol red  
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)  
(detection of H. pylori in stomach)

IT 14798-03-9, Ammonium, analysis  
RL: ARU (Analytical role, unclassified); FMU (Formation, unclassified); ANST (Analytical study); FORM (Formation, nonpreparative)  
(detection of H. pylori in stomach)


IT 9002-13-5, Urease  
RL: ANT (Analyte); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
(detection of H. pylori in stomach)

RN 9002-13-5 HCAPLUS  
CN Urease (8CI, 9CI) (CA INDEX NAME)


\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

IT 57-13-6, Urea, uses 143-74-8, Phenol red  
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)  
(detection of H. pylori in stomach)

RN 57-13-6 HCAPLUS  
CN Urea (8CI, 9CI) (CA INDEX NAME)



RN 143-74-8 HCAPLUS  
CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis- (9CI) (CA INDEX NAME)



- L69 ANSWER 8 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
 AN 2001:539704 HCAPLUS  
 DN 136:18612  
 TI One hundred years of discovery and rediscovery of *Helicobacter pylori* and its association with peptic ulcer disease  
 AU **Marshall, Barry J.**  
 CS Department of Microbiology QEII Medical Centre, University of Western Australia, Nedlands, 6009, Australia  
 SO *Helicobacter pylori* (2001), 19-24. Editor(s): Mobley, Harry L. T.; Mendz, George L.; Hazell, Stuart L. Publisher: ASM Press, Herndon, Va.  
 CODEN: 69BOCI  
 DT Conference; General Review  
 LA English  
 CC 14-0 (Mammalian Pathological Biochemistry)  
 AB A review of major highlights during the 100 yr of study of *Helicobacter pylori*. Topics discussed include spiral bacteria, epidemic **gastritis** with hypochlorhydria, the origin of **gastric urease**, and bismuth salts for **gastric** disease.  
 ST review *Helicobacter pylori* peptic ulcer  
 IT *Helicobacter pylori*  
 Stomach  
 (discovery and rediscovery of *Helicobacter pylori* and its assocn. with peptic ulcer disease)  
 IT Ulcer  
 (peptic; discovery and rediscovery of *Helicobacter pylori* and its assocn. with peptic ulcer disease)  
 IT **9002-13-5, Urease**  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (discovery and rediscovery of *Helicobacter pylori* and its assocn. with peptic ulcer disease)  
 IT 7440-69-9D, Bismuth, salts  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (discovery and rediscovery of *Helicobacter pylori* and its assocn. with peptic ulcer disease)  
 RE.CNT 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE  
 (1) Anon; The Principles and Practice of Medicine, 9th ed 1920  
 (2) Bizzozzero, G; Arch Mikr Anat 1893, V42, P82  
 (3) Blaser, M; J Physiol Pharmacol 1997, V48, P307 MEDLINE  
 (4) Blaser, M; Lancet 1997, V349, P1020 MEDLINE  
 (5) Castiglioni, A; A History of Medicine, 2nd ed 1947  
 (6) Chen, M; Lancet 1995, V339, P1120  
 (7) Coghlan, J; Lancet 1987, Vii, P1109  
 (8) Doenges, J; Arch Pathol 1939, V27, P469  
 (9) Ericsson, C; Rev Infect Dis 1986, V8(Suppl 2), PS202  
 (10) Figura, N; *Helicobacter* 1996, V1, P4 MEDLINE

- (11) Fitzgerald, O; Ir J Med Sci 1950, V292, P97
- (12) Freedberg, A; Am J Dig Dis 1940, V7, P443
- (13) Gibbons, A; Gastroenterology 1997, V112, P1940 HCAPLUS
- (14) Gledhill, T; Br Med J Clin Res Ed 1985, V290, P1383 MEDLINE
- (15) Graham, D; Ann Intern Med 1991, V115, P266 MEDLINE
- (16) Graham, D; Lancet 1987, Vi, P1174
- (17) Gregory, M; S Afr Med J 1982, V62, P52 MEDLINE
- (18) Harford, W; Gut, in press
- (19) Heilmann, K; Gut 1991, V32, P137 MEDLINE
- (20) Hirschowitz, B; Lancet 1956, Vii, P1081
- (21) Ito, S; Alimentary Canal 1967, P705
- (22) Krienitz, W; Dtsch Med Wochenschr 1906, V32, P872
- (23) Lambert, J; Med J Aust (Letter) 1985, V143, P174 MEDLINE
- (24) Langenberg, M; Lancet 1984, Vi, P1348
- (25) Lee, A; Infect Immun 1988, V56, P2843 MEDLINE
- (26) Lockard, V; Am J Vet Res 1970, V31, P1453 MEDLINE
- (27) Luck, J; Biochem J 1924, V18, P1227 HCAPLUS
- (28) Marshall, B; Am J Gastroenterol 1987, V82, P200 MEDLINE
- (29) Marshall, B; Digestion 1987, V37(Suppl 2), P16
- (30) Marshall, B; J Nucl Med 1988, V29, P11 MEDLINE
- (31) Marshall, B; Lancet 1986, Vi, P965
- (32) Marshall, B; Med J Aust 1985, V142, P436 MEDLINE
- (33) Matysiak-Budnik, T; Lancet (Letter) 1995, V346, P1489 MEDLINE
- (34) McNulty, C; Lancet 1984, Vi, P1068
- (35) Morris, A; Ann Intern Med 1991, V114, P662 MEDLINE
- (36) Ogle, J; Br Med J 1964, V1, P249
- (37) Palmer, E; Gastroenterology 1954, V27, P218
- (38) Peterson, W; Am J Gastroenterol 1993, V88, P2038 MEDLINE
- (39) Ramsey, E; Gastroenterology 1979, V76, P1449 MEDLINE
- (40) Rauws, E; Lancet 1990, V335, P1233 MEDLINE
- (41) Rotter, J; N Engl J Med 1979, V300, P63 MEDLINE
- (42) Salomon, H; Zentralbl Bakteriell 1896, V19, P433
- (43) Steer, H; Gut 1975, V16, P590 MEDLINE
- (44) Steer, H; Gut 1984, V25, P1203 MEDLINE
- (45) Sumner, J; J Biol Chem 1926, V69, P435 HCAPLUS
- (46) Takata, T; Gastrointest Endosc 1998, V47, P291 MEDLINE
- (47) Warren, J; Lancet 1983, Vi, P1273
- (48) Watanabe, T; Gastroenterology 1998, V115, P642 MEDLINE
- (49) Wiersinga, W; Gastroenterology 1977, V73, P1413 MEDLINE

IT 9002-13-5, Urease

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(discovery and rediscovery of Helicobacter pylori and its assocn. with  
peptic ulcer disease)

RN 9002-13-5 HCAPLUS

CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 9 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 2000:594938 HCAPLUS

DN 134:82869

TI A test strip for the estimation of urea in serum

AU Kumar, Hemant; Kumar, Ashok; Kumari, Poonam; Tulsani, N. B.

CS Centre for Biochemical Technology, Delhi, 110007, India

SO Indian Journal of Clinical Biochemistry (2000), 15(2), 124-127

CODEN: IJCBEY; ISSN: 0970-1915

PB Association of Clinical Biochemists of India

DT Journal

LA English

CC 9-2 (Biochemical Methods)

AB We have developed a biostrip for detn. of urea in serum. The  
test strip is based on enzymic assay where urease has been  
immobilized on the chromatog. paper along with chromogen,



**phenol red.** The **chromogen** is easily sol. in water and does not require other components for the color change. Serum **urea** reacts with **urease** and water to liberate **ammonia** and carbon dioxide. The liberated **ammonia** changes the pH of the reaction medium, which is monitored by the **chromogen phenol red**. A single step working reagent strip has been developed and the reaction is completed within 50 s at room temp. With this test strip **urea** concn. is measured in serum as low as 0.15 g/L. The speed and convenience of detg. **urea** in serum by this strip instantly makes it well suited for individuals, physicians and emergency centers.

ST **urea** detn serum biostrip enzymic

IT Blood analysis

(**urea** detn. in serum using biostrip)

IT 57-13-6, **Urea**, analysis

RL: ANT (Analyte); ANST (Analytical study)

(**urea** detn. in serum using biostrip)

RE.CNT 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Assadi, F; J Pediatr 1986, V108, P995 MEDLINE

(2) Court, J; Med J Aust 1972, V1, P525 MEDLINE

(3) Hurlburt, T; Amer J Clin Pathol 1991, V96(5), P582

(4) Jain, A; J AOAC Int 1999, V82(1), P9 HCAPLUS

(5) Lequang, N; Clin Chem 1987, V33, P192 MEDLINE

(6) Lespinas, F; Clin Chem 1989, V35(4), P654 HCAPLUS

(7) Li, X; Application and preparation of a BUN test strip 1993, V15(2), P152 HCAPLUS

HCAPLUS

(8) Marshall, S; Clin Chem 1992, V38, P588 HCAPLUS

(9) Naslund, B; Clin Chem 1998, V44, P1964 HCAPLUS

(10) Orsonneau, J; Clin Chem 1992, V38, P619 HCAPLUS

(11) Rogge, J; Amer J Gastroenterol 1995, V90(11), P1965 MEDLINE

(12) Tarnoky, A; Clin Chem Acta 1964, V10, P253 MEDLINE

(13) Waybenga, D; Clin Chem 1971, V17, P891


IT 57-13-6, **Urea**, analysis

RL: ANT (Analyte); ANST (Analytical study)

(**urea** detn. in serum using biostrip)

RN 57-13-6 HCAPLUS

CN Urea (8CI, 9CI) (CA INDEX NAME)



L69 ANSWER 10 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 2000:464644 HCAPLUS

DN 133:86488

TI An accurate and inexpensive method for measuring **urea** nitrogen using **urease**

IN Fujii, Takayuki

PA Yatron Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 5 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

IC ICM C12Q001-58

ICS G01N033-62

CC 9-16 (Biochemical Methods)

FAN.CNT 1

PATENT NO.


KIND

DATE


APPLICATION NO.

DATE

PI JP 2000189196 A2 20000711 JP 1998-376480 19981225  
 PRAI JP 1998-376480 19981225  
 AB An accurate method is provided for measuring **urea** nitrogen in a wide range of concn. using an inexpensive reagent. The **urea** nitrogen is quantitated by optically measuring the **pH** change due to **ammonia** generated upon reacting **urease** with **urea** in the presence of a **pH indicator** in the liq. phase contg. at least two kinds of **buffer**. The reagent comprises the first reagent component consisting of at least two kinds of **buffer** contg. at least a **pH indicator**, and the second reagent component consisting of the soln. contg. at least **urease**. The combination of two **buffer** solns. (e.g., HEPES and Tricine, TAPSO and EPPS) gave a linear calibration curve with **urea** in a wide range of concn., comparing with the cases where only one **buffer** soln. was used.  
 ST **urea** nitrogen **urease** **pH indicator**  
 IT **Acid-base indicators**  
     **Buffers**  
     (accurate and inexpensive method for measuring **urea** nitrogen using **urease**)  
 IT Calibration  
     (linear; accurate and inexpensive method for measuring **urea** nitrogen using **urease**)  
 IT 57-13-6, **Urea**, analysis 7727-37-9, Nitrogen, analysis  
     14798-03-9, Ammonium, analysis  
     RL: ANT (Analyte); ANST (Analytical study)  
     (accurate and inexpensive method for measuring **urea** nitrogen using **urease**)  
 IT 143-74-8, **Phenol Red** 2411-89-4,  
     o-Cresolphthaleincomplexone 9002-13-5, **Urease**  
     RL: ARG (Analytical reagent use); ANST (Analytical study); USES  
     (Uses)  
     (accurate and inexpensive method for measuring **urea** nitrogen using **urease**)  
 IT 5704-04-1, Tricine 7365-45-9, HEPES 16052-06-5, EPPS 68399-81-5,  
     TAPSO  
     RL: ARU (Analytical role, unclassified); ANST (Analytical study)  
     (accurate and inexpensive method for measuring **urea** nitrogen using **urease**)  
 IT 57-13-6, **Urea**, analysis  
     RL: ANT (Analyte); ANST (Analytical study)  
     (accurate and inexpensive method for measuring **urea** nitrogen using **urease**)  
 RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)



IT 143-74-8, **Phenol Red** 9002-13-5,  
     **Urease**  
     RL: ARG (Analytical reagent use); ANST (Analytical study); USES  
     (Uses)  
     (accurate and inexpensive method for measuring **urea** nitrogen using **urease**)  
 RN 143-74-8 HCAPLUS  
 CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis- (9CI) (CA INDEX NAME)



RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 11 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 1999:748430 HCAPLUS.

DN 131:348779

TI An immunological method for detecting Helicobacter pylori

IN Nakamura, Michihiro

PA Nihon Kodan Kogyo Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 4 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

IC ICM C12Q001-04

ICS G01N033-531; G01N033-569; G01N033-573

CC 9-10 (Biochemical Methods)

Section cross-reference(s): 10, 14

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 11318490	A2	19991124	JP 1998-136256	19980519
PRAI	JP 1998-136256		19980519		

AB A simple immunol. method is described for detecting Helicobacter pylori by specifically detecting the **urease** derived from Helicobacter pylori without using an exclusive and particular app. A **sample** liq. is contacted with the solid phase on which antibodies specific to **urease** derived from Helicobacter pylori is immobilized. Then, the solid phase is washed with a **coloring** liq. contg. at least **urea** and a **pH indicator**. After the washing step, the solid phase is contacted with the **coloring** liq. A change in the **color** of the **coloring** agent is measured or detected with the naked eye. By this method, Helicobacter pylori **urease** was detected at the concn. of more than 0.1mIU/mL.

ST Helicobacter pylori **urease** immunoassay **pH indicator**

IT **Acid-base indicators**  
 Helicobacter pylori  
 Immunoassay

(immunol. method for detecting Helicobacter pylori)

IT Antibodies

RL: ARG (Analytical reagent use); PEP (Physical, engineering or chemical process); ANST (Analytical study); PROC (Process); USES (Uses)  
 (to Helicobacter pylori **urease**; immunol. method for detecting Helicobacter pylori)

IT 9002-13-5, Urease

RL: ANT (Analyte); THU (Therapeutic use); ANST (Analytical

**study**); BIOL (Biological study); USES (Uses)  
 (immunol. method for detecting Helicobacter pylori)

IT 57-13-6, Urea, analysis 143-74-8,  
**Phenol red**  
 RL: ARU (Analytical role, unclassified); ANST (Analytical study)  
 (immunol. method for detecting Helicobacter pylori)


IT 9002-13-5, Urease  
 RL: ANT (Analyte); THU (Therapeutic use); ANST (Analytical  
**study**); BIOL (Biological study); USES (Uses)  
 (immunol. method for detecting Helicobacter pylori)

RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)


\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

IT 57-13-6, Urea, analysis 143-74-8,  
**Phenol red**  
 RL: ARU (Analytical role, unclassified); ANST (Analytical study)  
 (immunol. method for detecting Helicobacter pylori)

RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)



RN 143-74-8 HCAPLUS  
 CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis- (9CI) (CA  
 INDEX NAME)



L69 ANSWER 12 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
 AN 1999:421809 HCAPLUS  
 DN 131:41823  
 TI Colorimetric assessment of the sensitivity of Helicobacter  
 pylori to antimicrobial substances  
 IN Zuccato, Alessandro  
 PA Consortia Laboratories, Italy  
 SO PCT Int. Appl., 26 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM C12Q001-18  
 ICS C12Q001-58; G01N033-62  
 CC 9-16 (Biochemical Methods)  
 Section cross-reference(s): 1, 7, 10  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9932656	A1	19990701	WO 1998-IT367	19981216
	W: AU, BA, BG, BR, CA, CN, CZ, HR, HU, ID, IL, IS, JP, LT, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	IT 1297510	B1	19991217	IT 1997-VR122	19971223
	AU 9917831	A1	19990712	AU 1999-17831	19981216
PRAI	IT 1997-VR122	A	19971223		
	WO 1998-IT367	W	19981216		
AB	A <b>colorimetric</b> method which is advantageously applicable to a culture medium of the liq. type, and which is further suitable for the evaluation in vitro of sensitivity and resistance of <i>Helicobacter pylori</i> to antimicrobial pharmaceuticals, is based on the <b>colorimetric</b> detection of bacterial growth of <i>Helicobacter pylori</i> stemming from an increase in bacterial <b>urease</b> concn., said <b>color</b> variation being made possible by a <b>pH color indicator</b> injected into the culture medium. A <b>colorimetric</b> kit for the assessment of <i>Helicobacter pylori</i> 's sensitivity and/or resistance to antimicrobial pharmaceuticals, said assessment being carried out with the naked eye and/or by spectrophotometric means, comprises: (a) a plurality of microwells and/or vessels made of transparent material and contg. predetd. antimicrobial substances and at suitably predetd. concns., said vessels being advantageously assembled in printed modules and further being packed in a sterile manner; (b) a culture medium for <i>Helicobacter pylori</i> contg. <b>urea</b> and a <b>color pH indicator</b> .				
ST	<b>colorimetric</b> <i>Helicobacter pylori</i> antimicrobial substances				
IT	<b>Acid-base indicators</b>				
	Antimicrobial agents				
	<b>Colorimetry</b>				
	Culture media				
	Growth, microbial				
	<i>Helicobacter pylori</i>				
	Hybridoma				
	Immunoassay				
	Test kits				
	UV and visible spectroscopy				
	(colorimetric assessment of sensitivity of <i>Helicobacter pylori</i> to antimicrobial substances)				
IT	Antibodies				
	RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses) (monoclonal; <b>colorimetric</b> assessment of sensitivity of <i>Helicobacter pylori</i> to antimicrobial substances)				
IT	<b>9002-13-5, Urease</b>				
	RL: ANT (Analyte); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); USES (Uses) (colorimetric assessment of sensitivity of <i>Helicobacter pylori</i> to antimicrobial substances)				
IT	<b>57-13-6, Urea</b> , biological studies <b>143-74-8, Phenol red</b>				
	RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses) (colorimetric assessment of sensitivity of <i>Helicobacter pylori</i> to antimicrobial substances)				
RE.CNT	9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD				
RE					
	(1) Behringwerke, A; DE 19547708 A 1997 HCAPLUS				
	(2) Dimotech Ltd; EP 0689842 A 1996 HCAPLUS				
	(3) Jackson, F; US 5439801 A 1995 HCAPLUS				
	(4) Kang, J; Abstracts of the General Meeting of the American Society for Microbiology 1997, V97(0), P146				

- (5) King Wing; US 5498528 A 1996 HCAPLUS  
 (6) Marshall, B; US 4748113 A 1988 HCAPLUS  
 (7) Mirshahi, F; Journal of Clinical Pathology 1998, V51(3), P250  
 (8) Roda, A; Analytical Biochemistry 1998, V264(1), P47 HCAPLUS  
 (9) Vasquez, A; Journal of Clinical Microbiology 1996, V34(5), P1232 HCAPLUS

IT 9002-13-5, Urease  
 RL: ANT (Analyte); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (colorimetric assessment of sensitivity of Helicobacter pylori to antimicrobial substances)

RN 9002-13-5 HCAPLUS

CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*


IT 57-13-6, Urea, biological studies 143-74-8,

Phenol red

RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (colorimetric assessment of sensitivity of Helicobacter pylori to antimicrobial substances)


RN 57-13-6 HCAPLUS

CN Urea (8CI, 9CI) (CA INDEX NAME)



RN 143-74-8 HCAPLUS

CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis- (9CI) (CA INDEX NAME)



L69 ANSWER 13 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 1998:754811 HCAPLUS

DN 130:94588

TI A quick, simple and reliable method for detection of urea in adulterated milk

AU Kumar, Ashok; Kumar, Hemant; Tulsani, N. B.; Joshi, A. P.

CS Centre for Biochemical Technology, Delhi, 110 007, India

SO Oriental Journal of Chemistry (1998), 14(2), 189-192

CODEN: OJCHEG; ISSN: 0970-020X

PB Oriental Scientific Publishing Co.

DT Journal

LA English

CC 17-1 (Food and Feed Chemistry)

AB Milk is an essential nutrient for human and animals. It is used as milk

as well as in the form of milk products. The main constituents of milk are carbohydrates, proteins, vitamins, minerals and water. Due to the large demand of milk, the milkmaids add water to it to make it more profitable. However, addn. of water results in the decrease of sp. gr. of the milk. For the same reason, an alternate route of adding **urea**, oils and detergents are found which not only maintains the sp. gr. of the milk but are also economical. This practice of adulterating the milk, though harmful for the human beings, is increasing day-by-day, particularly in metropolis. Therefore, a quick and reliable method is required to detect the **urea** in the milk by the customers as well as by the supervising inspectors. A quick, simple, reliable and economical method for the detection of **urea** in the milk was developed. This involves the use of **urease**, which reacts with the milk **urea** to liberate **ammonia**. Subsequently, the liberated **ammonia** reacts with a specific dye and the color of the milk is changed to a blue color. The development of color is quick and is visible with the naked eye. The control will not show any color even after adding the dye.

ST **urea detection milk color indicator**

IT **Colorimetric indicators**

Milk analysis

(a quick, simple and reliable method for detection of **urea** in adulterated milk)

IT **57-13-6, Urea, analysis**

RL: ANT (Analyte); POL (Pollutant); ANST (Analytical study); OCCU (Occurrence)

(a quick, simple and reliable method for detection of **urea** in adulterated milk)

IT **76-59-5, Bromothymol blue 9002-13-5, Urease**

RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)

(a quick, simple and reliable method for detection of **urea** in adulterated milk)

RE.CNT 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Anon; ISI Handbook of Food Analysis V17(Part XII)
- (2) Anon; Official Methods of Analysis of AOAC International 16th Ed V2, P33
- (3) Arora, K; Indian Dairy man 1995, V47, P47
- (4) Arora, N; J Food Sci Technol 1986, V23, P213 HCAPLUS
- (5) Bansal, P; Current Science 1997, V73(11), P904
- (6) Chaney, A; Clin Chem 1962, V8, P130 HCAPLUS
- (7) Choudhary, M; Dudh ka Dudh Pani ka Pani Kaun Karega Navbharat Times 1997
- (8) Fazzio, T; Official methods of analysis of AOAC International 16th Ed V19(2), P47
- (9) Harrison, G; Chemical Methods in Clinical Medicine 3rd Ed 1947
- (10) Koch, F; J Amer Chem Soc 1924, V46, P2006
- (11) Malhotra, O; Indian J Biochem Biophysics 1969, V6, P15 HCAPLUS
- (12) Mathew, M; The Hindustan Times 1997
- (13) Sharma, S; The Hindustan Times 1996
- (14) Singhal, O; NDRI Annual Report 1993, V81
- (15) VanSlyke, D; J Biol Chem 1914, V19, P21


IT **57-13-6, Urea, analysis**

RL: ANT (Analyte); POL (Pollutant); ANST (Analytical study); OCCU (Occurrence)

(a quick, simple and reliable method for detection of **urea** in adulterated milk)

RN **57-13-6 HCAPLUS**

CN **Urea (8CI, 9CI) (CA INDEX NAME)**



IT 9002-13-5, Urease  
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES  
 (Uses)  
 (a quick, simple and reliable method for detection of urea in  
 adulterated milk)  
 RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 14 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 1998:711352 HCAPLUS

DN 130:48979

TI Development of a chemiluminescent **urease** activity assay for  
 Helicobacter pylori infection diagnosis in **gastric** mucosa  
**biopsies**

AU Roda, Aldo; Piazza, Francesco; Pasini, Patrizia; Baraldini, Mario;  
 Zambonin, Laura; Fossi, Stefania; Bazzoli, Franco; Roda, Enrico  
 CS Department of Pharmaceutical Sciences, University of Bologna, Bologna,  
 Italy

SO Analytical Biochemistry (1998), 264(1), 47-52  
 CODEN: ANBCA2; ISSN: 0003-2697

PB Academic Press

DT Journal

LA English

CC 7-1 (Enzymes)

Section cross-reference(s): 1, 9, 14

AB A chemiluminescent **urease** activity assay has been developed and  
 optimized using the chemiluminescent **pH indicator**  
 phthalhydrazidylazoacetylacetone. This compd. is stable at **pH**  
 .ltoreq. 7 and decomp. at higher **pH** values, emitting light in  
 the presence of H2O2. **Urease** catalyzes hydrolysis of  
**urea** to form **NH3** and **CO2** which increase the **pH**  
 of the reaction medium, thus allowing the chemiluminescent  
**indicator** to decomp. and produce photons. The emitted light is  
 proportional to the **urease** activity when **urea** is in  
 excess. **Urease** tests based on **colorimetric pH**  
**indicators** like **phenol red** are com. available  
 and commonly used for the rapid diagnosis of Helicobacter pylori infection  
 in **gastric** mucosa **biopsy** specimens, since this  
 bacterium produces high amts. of **urease**. Such  
**colorimetric** tests often lack sensitivity, giving false-neg.  
 results. The developed chemiluminescent test proved to be at least  
 50-fold more sensitive than the **colorimetric** tests, permitting  
 early diagnosis of infection, and it is more rapid, giving results in 1-10  
 min compared to 30 min. Further applications of this assay could be the  
 in situ localization of **urease** activity, corresponding to the  
 presence of H. pylori, in **gastric** mucosa cryosections and the  
 development of high throughput screening assays of antimicrobial drugs  
 able to inactivate the bacterium. (c) 1998 Academic Press.

ST **urease** chemiluminescent assay Helicobacter **gastric**  
 mucosa infection phthalhydrazidylazoacetylacetone

IT Infection  
 (bacterial; development of a chemiluminescent **urease** activity  
 assay for Helicobacter pylori infection diagnosis in **gastric**  
 mucosa **biopsies**)

IT Bioassay



## Diagnosis

Helicobacter pylori

Luminescence, chemiluminescence

(development of a chemiluminescent **urease** activity assay for  
Helicobacter pylori infection diagnosis in **gastric** mucosa  
**biopsies**)

IT Stomach

(mucosa; development of a chemiluminescent **urease** activity  
assay for Helicobacter pylori infection diagnosis in **gastric**  
mucosa **biopsies**)

IT 9002-13-5, Urease

RL: **ANT (Analyte)**; BAC (Biological activity or effector, except  
adverse); BSU (Biological study, unclassified); THU (Therapeutic use);  
**ANST (Analytical study)**; BIOL (Biological study); USES (Uses)  
(development of a chemiluminescent **urease** activity assay for  
Helicobacter pylori infection diagnosis in **gastric** mucosa  
**biopsies**)

IT 109632-03-3P

RL: ARG (Analytical reagent use); SPN (Synthetic preparation); ANST  
(Analytical study); PREP (Preparation); USES (Uses)  
(development of a chemiluminescent **urease** activity assay for  
Helicobacter pylori infection diagnosis in **gastric** mucosa  
**biopsies**)

IT 123-54-6, Acetylacetone, reactions 521-31-3, Luminol

RL: RCT (Reactant); RACT (Reactant or reagent)  
(development of a chemiluminescent **urease** activity assay for  
Helicobacter pylori infection diagnosis in **gastric** mucosa  
**biopsies**)

RE.CNT 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE

- (1) Abdalla, S; J Clin Microbiol 1989, V27, P2604 MEDLINE
- (2) Acs Committee On Environmental Improvement; Anal Chem 1980, V52, P2242
- (3) Barthel, J; Rev Infect Dis 1990, V12, P107
- (4) Bazzoli, F; Eur J Gastroenterol Hepatol 1994, V6, P773
- (5) Bronstein, I; Anal Biochem 1989, V180, P95 HCAPLUS
- (6) Campbell, A; Chemiluminescence Principles and Applications in Biology and  
Medicine 1988, P414
- (7) Coudron, P; J Clin Microbiol 1989, V27, P1527 MEDLINE
- (8) Crabtree, J; Gut 1993, V34, P1339 MEDLINE
- (9) Crabtree, J; Helicobacter pylori: Techniques for Clinical Diagnosis and  
Basic Research 1996, P235
- (10) Daskapoulos, G; Am J Gastroenterol 1994, V89, P1350
- (11) Ernst, P; Helicobacter pylori 1994, P295
- (12) Fauchere, J; Helicobacter pylori: Techniques for Clinical Diagnosis and  
Basic Research 1996, P50
- (13) Graham, D; Ann Intern Med 1992, V116, P705 MEDLINE
- (14) Graham, D; Gastroenterology 1991, V100, P1495 MEDLINE
- (15) Graham, D; Lancet 1987, V1, P1174 MEDLINE
- (16) Hazell, S; Am J Gastroenterol 1987, V82, P292 MEDLINE
- (17) Hazell, S; Helicobacter pylori 1994, P85
- (18) Lamouliatte, H; Helicobacter pylori: Techniques for Clinical Diagnosis and  
Basic Research 1996, P1
- (19) Loffeld, R; J Clin Pathol 1988, V41, P85 MEDLINE
- (20) Loffeld, R; J Pathol 1991, V165, P69 MEDLINE
- (21) Logan, R; Eur J Gastroenterol Hepatol 1991, V3, P915
- (22) Logan, R; Gut 1991, V32, P1461 MEDLINE
- (23) Logan, R; Helicobacter pylori: Techniques for Clinical Diagnosis and Basic  
Research 1996, P74
- (24) Marshall, B; Helicobacter pylori 1994, P75
- (25) Marshall, B; Lancet 1984, V1, P1311 MEDLINE
- (26) Megraud, F; J Clin Microbiol 1989, V27, P1870 MEDLINE
- (27) Mitchell, H; J Infect Dis 1992, V166, P149 MEDLINE
- (28) Musiani, M; Am J Pathol 1996, V148, P1105 HCAPLUS

- (29) Nomura, A; N Engl J Med 1991, V325, P1132 MEDLINE
- (30) Parsonnet, J; N Engl J Med 1991, V325, P1127 MEDLINE
- (31) Perez-Perez, G; Ann Intern Med 1988, V109, P11 MEDLINE
- (32) Peterson, W; N Engl J Med 1991, V324, P1043 MEDLINE
- (33) Potters, H; Histopathology 1987, V11, P1223 MEDLINE
- (34) Price, A; Helicobacter pylori: Techniques for Clinical Diagnosis and Basic Research 1996, P33
- (35) Rauws, E; Lancet 1990, V335, P1233 MEDLINE
- (36) Roda, A; Anal Biochem 1998, V257, P53 HCAPLUS
- (37) Roda, A; Anal Chem 1996, V68, P1073 HCAPLUS
- (38) Rokkas, T; Am J Gastroenterol 1987, V82, P1149 MEDLINE
- (39) Solte, N; Lancet 1992, V339, P745
- (40) Talley, N; J Clin Microbiol 1991, V29, P1635 MEDLINE
- (41) Thankarajan, N; J Indian Chem Soc 1986, V63, P977 HCAPLUS
- (42) Thankarajan, N; Talanta 1987, V34, P507 HCAPLUS
- (43) Westblom, T; J Clin Microbiol 1988, V26, P1393 MEDLINE
- (44) Wyatt, J; Histopathology 1995, V26, P1 MEDLINE

## IT 9002-13-5, Urease

RL: **ANT (Analyte)**; BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use);  
**ANST (Analytical study)**; BIOL (Biological study); USES (Uses)  
 (development of a chemiluminescent **urease** activity assay for  
 Helicobacter pylori infection diagnosis in **gastric** mucosa  
**biopsies**)

RN 9002-13-5 HCAPLUS

CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 15 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 1996:126652 HCAPLUS

DN 124:170019

TI Detection of Helicobacter pylori

IN King, Wing

PA USA

SO PCT Int. Appl., 35 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM C12Q001-04

ICS C12Q001-02; G01N033-53

CC 9-11 (Biochemical Methods)

Section cross-reference(s): 10, 14

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9534677	A1	19951221	WO 1995-US7598	19950612
	W: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN				
	RW: KE, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
	US 5498528	A	19960312	US 1994-257862	19940610
	AU 9528299	A1	19960105	AU 1995-28299	19950612
PRAI	US 1994-257862		19940610		
	WO 1995-US7598		19950612		

AB A method for detecting H. pylori is disclosed that involves contacting a **sample** suspected of contg. H. pylori with a medium which provides for substantially selective growth of H. pylori, incubating the **sample** with the medium for a time sufficient for detection of H. pylori growth and detecting the growth and thereby reducing the presence of H. pylori within the **sample**. The methodol. employs a wide

range of a different culture media which are modified specifically for the selective growth and specific detection of *H. pylori*. A typical medium includes Columbia broth supplemented with **urea** and a **pH indicator**. The methodol. provides for a relatively high degree of sensitivity (i.e., small nos. of bacteria present within a **sample** are detected) as well as high selectivity (i.e., the method provides for a low percentage of false positives). Various kits used in connection with the method are designed so that they can be used by unskilled users in an "at home" setting. The kits and methodol. are economical, easily used and provide highly accurate results within a relatively short period of time (e.g., 3 days or less).

- ST Helicobacter pylori detection culture media kit; stomach **biopsy**
- IT Helicobacter pylori detection; body fluid Helicobacter pylori detection
- IT Stomach
  - (**biopsy**; culture media compns. and methods and kits for Helicobacter pylori detection)
- IT Animal tissue
  - Blood analysis
  - Blood serum
  - Campylobacter pyloridis
  - Culture media
  - Feces
  - Microorganism growth
  - Saliva
  - Yeast
    - (culture media compns. and methods and kits for Helicobacter pylori detection)
- IT Antibodies
  - RL: ANT (Analyte); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)
  - (culture media compns. and methods and kits for Helicobacter pylori detection)
- IT Digestive tract
  - (secretion; culture media compns. and methods and kits for Helicobacter pylori detection)
- IT **Indicators**
  - (acid-base, culture media compns. and methods and kits for Helicobacter pylori detection)
- IT Caseins, biological studies
  - RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
  - (hydrolyzates, culture media compns. and methods and kits for Helicobacter pylori detection)
- IT **9002-13-5, Urease**
  - RL: **ANT (Analyte)**; CAT (Catalyst use); **ANST (Analytical study)**; USES (Uses)
  - (culture media compns. and methods and kits for Helicobacter pylori detection)
- IT 50-99-7, Glucose, biological studies 52-89-1, L-Cysteine hydrochloride 57-13-6, **Urea**, biological studies 68-04-2, Sodium citrate 77-86-1, Tris **buffer** 738-70-5, Trimethoprim 1404-90-6, Vancomycin 7487-88-9, Magnesium sulfate, biological studies 7647-14-5, Sodium chloride, biological studies 7720-78-7 12633-72-6, Amphotericin 103370-88-3, IsoVitaleX
  - RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
  - (culture media compns. and methods and kits for Helicobacter pylori detection)
- IT **9002-18-0, Agar**
  - RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
  - (culture media contg.; culture media compns. and methods and kits for Helicobacter pylori detection)


IT 9002-13-5, Urease  
 RL: ANT (Analyte); CAT (Catalyst use); ANST (Analytical study); USES (Uses)  
 (culture media compns. and methods and kits for Helicobacter pylori detection)

RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

IT 57-13-6, Urea, biological studies  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
 (culture media compns. and methods and kits for Helicobacter pylori detection)

RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)



IT 9002-18-0, Agar  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
 (culture media contg.; culture media compns. and methods and kits for Helicobacter pylori detection)

RN 9002-18-0 HCAPLUS  
 CN Agar (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 16 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 1995:665321 HCAPLUS

DN 123:51695

TI Detection of Helicobacter pylori in the stomach using urea- and indicator-containing reagents

IN Marshall, Barry

PA USA

SO PCT Int. Appl., 16 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K009-28

ICS A61K009-48; A91K009-54; C12Q001-04; C12Q001-58; G01N021-77

CC 9-2 (Biochemical Methods)

Section cross-reference(s): 7

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9511672	A1	19950504	WO 1994-US12332	19941025
	W: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ				
	RW: KE, MW, SD, SZ, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
	CA 2174933	AA	19950504	CA 1994-2174933	19941025
	AU 9481270	A1	19950522	AU 1994-81270	19941025
	EP 725633	A1	19960814	EP 1995-900448	19941025
	R: AT, CH, DE, GB, IE, LI, LU				

CN 1139381	A	19970101	CN 1994-194624	19941025
JP 09506246	T2	19970624	JP 1994-512826	19941025
BR 9407718	A	19971111	BR 1994-7718	19941025

PRAI US 1993-142600 A 19931028  
WO 1994-US12332 W 19941025

AB A method for the in vivo detection of **urease**-producing helicobacter in the upper stomach is disclosed. The dense carrier is divided into two sep. groups which are combined with sep. reagent **indicators**, one of which also contains **urea**. The carriers are food sol. products, preferably sugar beads having a diam. of approx. 0.2 to 3.0 mm. The treated carriers and **urea** are encapsulated in a sol. capsule which is administered to a patient. The d. of the carriers cause the capsule to migrate to the **gastric** mucosa, where the capsule is dissolved, placing the reagents and **urea** in direct contact with the **gastric** mucosa. The **urea** reacts with any **urease** present in the stomach by creating **ammonia**, which increases the **pH** within the stomach. The two reagents react differently, through **color** change, to the increase in **pH**, which is viewed through use of an endoscope. A preferred first reagent is bromothymol blue (dibromothymolsulfonphthalein), which changes yellow in the presence of **urease**, and a preferred second reagent is **phenol red** (phenolsulfonphthalein) which turns red in the presence of **urease**.

ST Helicobacter detection stomach **urea indicator**;  
**urease** Helicobacter detection stomach; bromothymol blue  
Helicobacter detection stomach; **phenol red**  
Helicobacter detection stomach

IT Helicobacter  
**Indicators**  
Stomach  
(Helicobacter in vivo detection in stomach with **urea-** and **indicator**-contg. reagents)

IT Carbohydrates and Sugars, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(Helicobacter in vivo detection in stomach with **urea-** and **indicator**-contg. reagents)

IT Food  
(sol. food products; Helicobacter in vivo detection in stomach with **urea-** and **indicator**-contg. reagents)

IT Medical goods  
Optical instruments  
(endoscopes, Helicobacter in vivo detection in stomach with **urea-** and **indicator**-contg. reagents)

IT 57-13-6, **Urea**, uses 76-59-5, Bromothymol blue  
143-74-8, **Phenol red** 594-05-8, **Urea**  
-14C 58069-82-2, **Urea**-13C  
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)  
(Helicobacter in vivo detection in stomach with **urea-** and **indicator**-contg. reagents)


IT 7664-41-7, **Ammonia**, analysis  
RL: ARU (Analytical role, unclassified); ANST (Analytical study)  
(Helicobacter in vivo detection in stomach with **urea-** and **indicator**-contg. reagents)

IT 9002-13-5, **Urease**  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
(Helicobacter in vivo detection in stomach with **urea-** and **indicator**-contg. reagents)


IT 57-13-6, **Urea**, uses 143-74-8, **Phenol red**  
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)

(Helicobacter in vivo detection in stomach with **urea-** and **indicator-contg.** reagents)

RN 57-13-6 HCAPLUS  
CN Urea (8CI, 9CI) (CA INDEX NAME)



RN 143-74-8 HCAPLUS  
CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis- (9CI) (CA INDEX NAME)



IT 7664-41-7, **Ammonia**, analysis  
RL: ARU (Analytical role, unclassified); ANST (Analytical study)  
(Helicobacter in vivo detection in stomach with **urea-** and **indicator-contg.** reagents)  
RN 7664-41-7 HCAPLUS  
CN Ammonia (8CI, 9CI) (CA INDEX NAME)

NH<sub>3</sub>

IT 9002-13-5, **Urease**  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
(Helicobacter in vivo detection in stomach with **urea-** and **indicator-contg.** reagents)  
RN 9002-13-5 HCAPLUS  
CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 17 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
AN 1994:503066 HCAPLUS  
DN 121:103066  
TI Device for carrying out **urease** tests for combined antrum/corpus **biopsies** to diagnose **gastrointestinal** diseases  
IN Heckenmueller, Harald; Meyer, Hansjoerg  
PA Astra Chemicals GmbH, Germany  
SO PCT Int. Appl., 24 pp.  
CODEN: PIXXD2  
DT Patent  
LA German  
IC ICM C12Q001-58

ICS B01L003-00; A61B010-00

CC 7-1 (Enzymes)

Section cross-reference(s): 14

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9413830	A1	19940623	WO 1993-DE1085	19931112
	W: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, PL, RO, RU, SD, SE, SK, US				
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
	AU 9454175	A1	19940704	AU 1994-54175	19931112
	AU 672657	B2	19961010		
	EP 673434	A1	19950927	EP 1993-924519	19931112
	EP 673434	B1	19990728		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, MC, NL, SE				
	JP 08506890	T2	19960723	JP 1993-513636	19931112
	AT 182627	E	19990815	AT 1993-924519	19931112
	BR 9307593	A	19990831	BR 1993-7593	19931112
	ES 2133417	T3	19990916	ES 1993-924519	19931112
	PL 177482	B1	19991130	PL 1993-309284	19931112
	US 5679570	A	19971021	US 1995-446841	19950601
	FI 9502767	A	19950606	FI 1995-2767	19950606
	NO 9502254	A	19950607	NO 1995-2254	19950607
PRAI	DE 1992-9217130		19921208		
	WO 1993-DE1085		19931112		
AB	The title device for diagnosis of <b>gastrointestinal</b> diseases assocd. with the <b>urea</b> -degrading bacterium <i>Helicobacter pylori</i> ( <i>Campylobacter pylori</i> ) has a carrier plate, a schematic representation of the stomach on the plate, at least one opening in the plate at the locations corresponding to the corpus and the antrum in the schematic representation of the stomach, an evaluation scale for assessment of the <b>urease</b> test and an area for data on the patient and for clin. data. A <b>gelled</b> substrate mixt. contg. yeast ext., KH2PO4, <b>phenol red</b> , <b>agar-agar</b> , dextrose, <b>urea</b> , vitamins, and trace elements is described for <b>urease</b> detection with the device.				
ST	stomach <b>urease</b> detection <b>gastrointestinal</b> disease diagnosis; app <b>urease</b> detection stomach <b>biopsy</b>				
IT	Yeast (ext., in <b>urease</b> detection in stomach <b>biopsies</b> for disease diagnosis)				
IT	Campylobacter pyloridis ( <b>gastrointestinal</b> diseases assocd. with, diagnosis of, <b>urease</b> detection in stomach <b>biopsies</b> in)				
IT	Stomach, composition (antrum, <b>urease</b> detection in <b>biopsy</b> of, in <b>gastrointestinal</b> disease diagnosis, app. for)				
IT	Stomach, composition (corpus, <b>urease</b> detection in <b>biopsy</b> of, in <b>gastrointestinal</b> disease diagnosis, app. for)				
IT	Digestive tract (disease, diagnosis of, <b>urease</b> detection in stomach <b>biopsies</b> in, app. for)				
IT	9002-13-5, Urease RL: ANT (Analyte); ANST (Analytical study) (detection of, in stomach <b>biopsies</b> in <b>gastrointestinal</b> disease diagnosis, app. for)				
IT	9002-18-0, Agar-agar 10043-35-3, Boric acid, uses 10361-37-2, Barium chloride, uses 50-99-7, Dextrose, uses 57-13-6, Urea, uses 58-85-5, D-(+)-Biotin 59-43-8, Vitamin B1, uses 59-67-6, Nicotinic acid, uses 68-19-9, Vitamin B12 139-33-3, Titriplex III 143-74-8, Phenol red				

150-13-0, p-Aminobenzoic acid 524-36-7, Pyridoxamine dihydrochloride  
 867-81-2, Sodium D-pantothenate 1344-13-4, Tin chloride 7447-40-7,  
 Potassium chloride, uses 7447-41-8, Lithium chloride, uses 7631-95-0,  
 Sodium molybdate 7646-79-9, Cobalt chloride, uses 7646-85-7, Zinc  
 chloride, uses 7720-78-7, Iron sulfate 7758-02-3, Potassium bromide,  
 uses 7758-98-7, Copper sulfate, uses 7773-01-5, Manganese chloride  
 7778-77-0, Potassium phosphate (KH<sub>2</sub>PO<sub>4</sub>)

RL: ANST (Analytical study)  
 (in **urease** detection in stomach **biopsies** for  
 disease diagnosis)

IT 9002-13-5, **Urease**

RL: ANT (Analyte); ANST (Analytical study)  
 (detection of, in stomach **biopsies** in  
**gastrointestinal** disease diagnosis, app. for)

RN 9002-13-5 HCAPLUS

CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

IT 9002-18-0, **Agar-agar** 57-13-6,  
**Urea**, uses 143-74-8, **Phenol red**

RL: ANST (Analytical study)  
 (in **urease** detection in stomach **biopsies** for  
 disease diagnosis)


RN 9002-18-0 HCAPLUS

CN Agar (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*


RN 57-13-6 HCAPLUS

CN Urea (8CI, 9CI) (CA INDEX NAME)



RN 143-74-8 HCAPLUS

CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis- (9CI) (CA  
 INDEX NAME)



L69 ANSWER 18 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 1993:554890 HCAPLUS

DN 119:154890

TI Reagents for easy determination of **urea**

IN Tabata, Yasushi; Suzuki, Hiroshi; Oomori, Masayuki

PA Terumo Corp, Japan

SO Jpn. Kokai Tokkyo Koho, 5 pp.



CODEN: JKXXAF  
 DT Patent  
 LA Japanese  
 IC ICM C12Q001-58  
 CC 7-1 (Enzymes)  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 05161500	A2	19930629	JP 1991-352913	19911217
PRAI	JP 1991-352913		19911217		

AB **Urea** (I) in body fluids is visually detd. with reagents comprising **urease**, **buffer**, and .gtoreq.2 **pH indicator** having different **color-changing pH** region. The method is accurate, and it can det. the concn. of **urea** in the range of 0.0 to 2.5%. Use of Bromthymol blue, **Phenol Red**, Phenolphthalein, and Thymol blue in detn. of I was shown.

ST **urea** easy detn reagent **pH indicator**;  
**urease** visual detn **urea** body fluid

IT **Buffer** substances and systems  
 (in reagents for easy visual detn. of **urea** in body fluids)

IT Body fluid  
 (**urea** easy visual detn. in, reagents contg. **urea** and **buffer** and **pH indicators** for)

IT **Indicators**  
 (acid-base, in reagents for easy visual detn. of **urea** in body fluids)


IT **57-13-6, Urea**, properties  
 RL: PRP (Properties)  
 (easy visual detn. of, in body fluids, reagents contg. **urea** and **buffer** and **pH indicators** for)

IT 76-59-5, Bromthymol blue 76-61-9, Thymol blue 77-09-8, Phenolphthalein  
**143-74-8, Phenol Red 9002-13-5, Urease**  
 RL: **ANST (Analytical study)**  
 (in reagents for easy visual detn. of **urea** in body fluids)

IT **57-13-6, Urea**, properties  
 RL: PRP (Properties)  
 (easy visual detn. of, in body fluids, reagents contg. **urea** and **buffer** and **pH indicators** for)

RN 57-13-6 HCAPLUS


CN Urea (8CI, 9CI) (CA INDEX NAME)



IT **143-74-8, Phenol Red 9002-13-5, Urease**  
 RL: **ANST (Analytical study)**  
 (in reagents for easy visual detn. of **urea** in body fluids)

RN 143-74-8 HCAPLUS

CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis- (9CI) (CA INDEX NAME)



RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 19 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
 AN 1992:569859 HCAPLUS  
 DN 117:169859  
 TI Rapid determination of **urea** in milk  
 IN Rajamaki, Sinikka; Riihimaki, Anne Maria  
 PA Valio Meijerien Keskusosuusliike, Finland  
 SO Brit. UK Pat. Appl., 14 pp.  
 CODEN: BAXXDU

DT Patent  
 LA English  
 IC ICM G01N033-52  
 ICS C12Q001-58; G01N033-04  
 CC 17-1 (Food and Feed Chemistry)  
 Section cross-reference(s): 13, 18

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	GB 2250590	A1	19920610	GB 1991-25040	19911126
	FI 9005949	A	19920604	FI 1990-5949	19901203
	FI 88310	B	19930115		
	FI 88310	C	19930426		
	SE 9103370	A	19920604	SE 1991-3370	19911114
	SE 512818	C2	20000522		
	NO 9104711	A	19920604	NO 1991-4711	19911129
	DK 9101950	A	19920604	DK 1991-1950	19911202
PRAI	FI 1990-5949	A	19901203		

AB A method for detn. of **urea** in milk that can be used in the field is described. In the method the reaction zone is sep. from the **indicator** zone and the **indicator** reaction of the detn. must be conducted in a closed vessel. The reaction uses **urease** to degrade **urea** to ammonium salts and these are then broken down to free **NH3** by exposure to an alk. The released **NH3** then changes the **color** of an **indicator** zone that contains a **pH**-dependent dye. The **color** of the incubator strip after a fixed incubation period is used for a semi-quant. detn. of **urea**. Filter paper strips were impregnated with a mixt. **Phenol Red** Na salt and **Bromothymol Blue** Na salt **buffered** with  $\text{NaH}_2\text{PO}_4$  at 0.00025 M or 0.0023 M. These were placed in the two-part cover of a tightly closable 10 mL container with the two papers kept sepd. by a partition. **Urease** (60-250 units) was added to the container and 1 mL milk added. After 2.5 min min.  $\text{NaOH}$  0.1 M was added and the cover closed. The **colors** of the test strips after a further 2.5 min showed a relationship to the **urea**

content of the milk as detd. by a spectrophotometric method of the prior art. Milk **samples** could be sorted in to <20, 20-30, or >30 mg **urea**/100 mL categories using this test. The method of the invention correctly classified 86.4% of milk **samples** compared to the spectrophotometric method.

ST milk **urea** detn **urease**

IT Phosphates, uses

RL: USES (Uses)

(as **buffer** in rapid detn. of **urea** in milk using **urease** and alk. and **pH**-dependent test strips)

IT Apparatus

(biochem., test strips, **pH**-dependent, in rapid detn. of **urea** in milk using **urease** and alk.)

IT Milk analysis

(for **urea** detn., rapid assay using **urease** and alk. and **pH**-dependent test strips for)

IT Indicators

(**pH**-dependent, in rapid detn. of **urea** in milk using **urease** and alk.)

IT **Buffer** substances and systems

(phosphate, in rapid detn. of **urea** in milk using **urease** and alk. and **pH**-dependent test strips)

IT 34487-61-1, **Phenol red**, sodium salt

34722-90-2

RL: ANST (Analytical study)

(as **indicator** dye in rapid detn. of **urea** in milk using **urease** and alk.)

IT 57-13-6, **Urea**, biological studies

RL: BIOL (Biological study)

(in milk, rapid detn. of, **urease** and alk. and **pH**-dependent test strips in)

IT 9002-13-5, **Urease**

RL: ANST (Analytical study)

(**urea** assay for milk using alk. and **pH**-dependent test strips and)


IT 34487-61-1, **Phenol red**, sodium salt

RL: ANST (Analytical study)

(as **indicator** dye in rapid detn. of **urea** in milk using **urease** and alk.)

RN 34487-61-1 HCAPLUS

CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis-, monosodium salt (9CI) (CA INDEX NAME)




Na

IT 57-13-6, **Urea**, biological studies

RL: BIOL (Biological study)  
(in milk, rapid detn. of, **urease** and alk. and pH  
-dependent test strips in)

RN 57-13-6 HCAPLUS  
CN Urea (8CI, 9CI) (CA INDEX NAME)



IT 9002-13-5, Urease  
RL: ANST (Analytical study)  
(urea assay for milk using alk. and pH-dependent  
test strips and)  
RN 9002-13-5 HCAPLUS  
CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*


L69 ANSWER 20 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
AN 1992:250813 HCAPLUS  
DN 116:250813  
TI New plate medium for growth and detection of **urease** activity of  
Helicobacter pylori  
AU Cellini, L.; Allocati, N.; Piccolomini, R.; Di Campi, E.; Dainelli, B.  
CS Fac. Med. Chir., Univ. "G. D'Annunzio", Chieti, 66013, Italy  
SO Journal of Clinical Microbiology (1992), 30(5), 1351-3  
CODEN: JCMIDW; ISSN: 0095-1137  
DT Journal  
LA English  
CC 7-1 (Enzymes)  
Section cross-reference(s): 10  
AB A new medium for detection of **urease** activity and isolation of  
H. pylori is proposed. This medium, contg. Columbia **Agar** Base,  
was supplemented with IsoVitaleX, hemin, **urea**, and  
**Phenol Red** (nonselective medium [NSM]). Both bacterial  
growth and **color** change were evaluated and compared with growth  
in the same medium supplemented with cefsulodin, vancomycin, polymyxin B  
sulfate, and amphotericin B (selective medium [SM]). Twenty-five recent  
clin. isolates and antral **biopsy** specimens from 33 patients who  
underwent endoscopy were examd. The isolates showed a rapid **color**  
change and good growth at 5 days of incubation with NSM and SM. H.  
pylori-pos. **biopsies** revealed a **color** change within 36  
h, and bacterial growth was better appreciated in NSM, but with more  
contaminating flora than in SM.  
ST culture medium Helicobacter isolation detection; **urease**  
detection Helicobacter culture medium  
IT Campylobacter pyloridis  
(detection and isolation of, culture medium for)  
IT 9002-13-5, Urease  
RL: ANT (Analyte); ANST (Analytical study)  
(detection of, in Helicobacter pylori, culture medium for)  
IT 9002-13-5, Urease  
RL: ANT (Analyte); ANST (Analytical study)  
(detection of, in Helicobacter pylori, culture medium for)  
RN 9002-13-5 HCAPLUS  
CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 21 OF 34 HCAPLUS COPYRIGHT 2003 ACS


AN 1991:509390 HCAPLUS  
 DN 115:109390  
 TI Diagnostic unit dose for the determination of **urease**  
 IN Rothgang, Gerhart; Mann, Helmut Josef; Klein, Cornelia J.  
 PA Roehm Pharma G.m.b.H., Germany  
 SO Eur. Pat. Appl., 7 pp.  
 CODEN: EPXXDW  
 DT Patent  
 LA German  
 IC ICM C12Q001-58  
 CC 7-1 (Enzymes)  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 369292	A1	19900523	EP 1989-120602	19891107
	R: AT, BE, CH, DE, ES, FR, GB, IT, LI, NL, SE				
PRAI	DE 1988-8814264		19881115		
AB	A convenient, dry solid unit dose of <b>urea</b> for detn. of <b>urease</b> contains <b>urea</b> (10-320 mg), a <b>buffer</b> for the <b>pH</b> range 5.0-7.5 (0.01-1 mg), a <b>pH indicator</b> which changes <b>color</b> over the <b>pH</b> range 5.5-8.5 (0.001-0.05 mg), and optionally a preservative (no data). The <b>buffer</b> may be KH <sub>2</sub> PO <sub>4</sub> -Na <sub>2</sub> HPO <sub>4</sub> , and the <b>indicator</b> may be e.g. bromcresol purple, neutral red, etc. The components may be compressed to a tablet.				
ST	<b>urease</b> detn <b>urea</b> reagent				
IT	<b>Buffer</b> substances and systems (solid units dose contg., for <b>urease</b> detn.)				
IT	<b>Indicators</b> (acid-base, solid unit dose contg., for <b>urease</b> detn.)				
IT	9002-13-5, <b>Urease</b> RL: <b>ANT (Analyte)</b> ; <b>ANST (Analytical study)</b> (detn. of, solid unit dose for)				
IT	57-13-6, <b>Urea</b> , biological studies RL: <b>BIOL (Biological study)</b> (solid units dose contg., for <b>urease</b> detn.)				
IT	9002-13-5, <b>Urease</b> RL: <b>ANT (Analyte)</b> ; <b>ANST (Analytical study)</b> (detn. of, solid unit dose for)				
RN	9002-13-5 HCAPLUS				
CN	Urease (8CI, 9CI) (CA INDEX NAME)				
*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***					
IT	57-13-6, <b>Urea</b> , biological studies RL: <b>BIOL (Biological study)</b> (solid units dose contg., for <b>urease</b> detn.)				
RN	57-13-6 HCAPLUS				
CN	Urea (8CI, 9CI) (CA INDEX NAME)				



L69 ANSWER 22 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
 AN 1991:3292 HCAPLUS  
 DN 114:3292  
 TI **Urea** protects *Helicobacter (Campylobacter) pylori* from the bactericidal effect of acid  
 AU **Marshall, B. J.**; **Barrett, L. J.**; **Prakash, C.**; **McCallum, R. W.**; **Guerrant, R. L.**

CS Dep. Med., Univ. Virginia, Charlottesville, VA, 22908, USA  
 SO Gastroenterology (1990), 99(3), 697-702  
 CODEN: GASTAB; ISSN: 0016-5085  
 DT Journal  
 LA English  
 CC 10-5 (Microbial Biochemistry)  
 AB Colonization of the stomach with *H. pylori* is common in patients with duodenal ulcers, which is known for its high acid secretion. Although the bacterium is usually isolated by culture of a **gastric biopsy** specimen, viable organisms may sometimes be found in the acidic **gastric** juice. It was postulated that **urease**, by generating **NH<sub>3</sub>**, protected *H. pylori* from acid. To test this hypothesis, the **pH** susceptibility of *H. pylori*, *Proteus mirabilis*, and the **urease-neg.** *Campylobacter jejuni* was examd. in the presence and absence of **urea**. It was found that without **urea** the 3 bacteria were all highly susceptible to acid. In striking contrast, the addn. of 5 mM **urea** completely protected *H. pylori*, but not *P. mirabilis* or *C. jejuni*, from **pH** values .gtoreq.1.5. The protective effect of **urea** on *H. pylori* was found with **urea** concns. .gtoreq.0.05 mM. The high **urease** activity of *H. pylori* apparently enables it to survive in **gastric** acid.  
 ST acid protection **urea urease** *Helicobacter*  
 IT *Campylobacter pyloridis*  
 (gastric acid effect on, **urea** protection against)  
 IT 57-13-6, **Urea**, biological studies  
 RL: BIOL (Biological study)  
 (in protection of *Helicobacter pylori* against **gastric** acid)  
 IT 9002-13-5, **Urease**  
 RL: BIOL (Biological study)  
 (of *Helicobacter pylori*, protection against **gastric** acid by)  
 IT 57-13-6, **Urea**, biological studies  
 RL: BIOL (Biological study)  
 (in protection of *Helicobacter pylori* against **gastric** acid)  
 RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)



IT 9002-13-5, **Urease**  
 RL: BIOL (Biological study)  
 (of *Helicobacter pylori*, protection against **gastric** acid by)  
 RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 23 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
 AN 1990:548589 HCAPLUS  
 DN 113:148589  
 TI **pH** regulation of **urease** levels in *Streptococcus salivarius*  
 AU Sissons, C. H.; Perinpanayagam, H. E. R.; Hancock, E. M.; Cutress, T. W.  
 CS Dent. Res. Unit, Med. Res. Counc. New Zealand, Wellington, N. Z.  
 SO Journal of Dental Research (1990), 69(5), 1131-7  
 CODEN: JDREAF; ISSN: 0022-0345  
 DT Journal  
 LA English  
 CC 10-2 (Microbial Biochemistry)

AB Potential mechanisms for regulation of **urease** levels in *S. salivarius* were examd., including induction by **urea**, N or C source repression, and effects of **pH** and CO<sub>2</sub> (because CO<sub>2</sub> enrichment enhanced **urease** detection on **urea agar** plates). Regulation by either **pH** or CO<sub>2</sub> was confirmed by comparison of the **urease** accumulation pattern during anaerobic growth under CO<sub>2</sub> with that under N<sub>2</sub>. Under CO<sub>2</sub>, there was an initial **buffering** plateau at **pH** 6.2 and a rate of *S. salivarius* **urease** accumulation 3-fold that under N<sub>2</sub>, with a **pH** 7.6 plateau. With both gas phases there was also an increase in the rate of **urease** appearance coincident with the decrease in medium **pH** following the **pH** plateau. The effects of **pH**, CO<sub>2</sub>, and HCO<sub>3</sub><sup>-</sup> on **urease** levels and on growth were sep. assessed by culture in media contg. 0, 25, or 100 mM KHCO<sub>3</sub> **buffered** at different **pH** levels. There was an inverse relation between the logarithm of the **urease** level after 24 h growth and the **pH** during growth; the **urease** sp. activity was 100-fold higher at **pH** 5.5 than at **pH** 7.0. HCO<sub>3</sub><sup>-</sup>/CO<sub>2</sub> (100 mM) had little effect on **urease** levels but was essential for growth at **pH** 5.5. There was no significant **urease** induction by **urea** or repression by NH<sub>3</sub> or glucose. There was also evidence of **pH** regulation of **urease** levels in some staphylococci, *Klebsiella pneumoniae*, and *Corynebacterium renale*, but not in *Actinomyces naeslundii* or several other species. Thus the external **pH** is a major factor regulating **urease** levels in *S. salivarius* and possibly some other species, a mechanism equiv. to **urease** repression by OH<sup>-</sup>.

ST Streptococcus **urease pH** carbon dioxide

IT Streptococcus salivarius  
(**urease** of, **pH** regulation of)

IT 9002-13-5, **Urease**

RL: PROC (Process)

(of Streptococcus salivarius, **pH** regulation of)

IT 71-52-3, Bicarbonate, biological studies 124-38-9, Carbon dioxide, biological studies

RL: BIOL (Biological study)

(**urease** of Streptococcus salivarius regulation by)

IT 9002-13-5, **Urease**

RL: PROC (Process)

(of Streptococcus salivarius, **pH** regulation of)

RN 9002-13-5 HCAPLUS

CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 24 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 1990:548382 HCAPLUS

DN 113:148382

TI Compositions and methods for the enrichment and isolation of *Campylobacter pylori* and related organisms

IN **Marshall, Barry J.**; Guerrant, Richard L.

PA University of Virginia Alumni Patents Foundation, USA

SO U.S., 3 pp.

CODEN: USXXAM

DT Patent

LA English

IC ICM C12Q001-58

ICS C12Q001-04; C12Q001-34; C12Q001-24

NCL 435012000

CC 9-2 (Biochemical Methods)

Section cross-reference(s): 10

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4923801	A	19900508	US 1987-37938	19870413
PRAI	US 1987-37938		19870413		
AB	C. pylori is enriched and isolated from a specimen contaminated with other organisms by (a) homogenizing the specimen (e.g. <b>gastric biopsy</b> , stool) with water; (b) introducing the specimen into a soln. of <b>urea</b> at pH 1.6 to req. 2.5 to kill nonurease-producing and some <b>urease</b> -producing organisms and to destroy preformed extracellular <b>urease</b> ; (c) plating the remaining <b>urease</b> -producing organisms onto a medium which contains antibiotics inhibitory to most of the remaining <b>urease</b> -producing organisms, but not inhibitory to C. pylori, and (d) detecting colonies of C. pylori. Stool inoculated with C. pylori was homogenized with saline and then mixed with 5 mM <b>urea</b> acidified to pH 1.6 with H2SO4. After 5 min at room temp., the specimen was plated onto nonselective blood <b>agar</b> and cultured for 3 days. After 3 days there were colonies of C. pylori and very few contaminating organisms on the plate.				
ST	Campylobacter isolation acid <b>urea</b>				
IT	Antibiotics (in Campylobacter pylori enrichment and isolation with <b>urea</b> and acids)				
IT	Campylobacter pyloridis (isolation of, <b>urea</b> and acids in)				
IT	Acids, biological studies RL: BIOL (Biological study) (Campylobacter pylori enrichment and isolation in presence of <b>urea</b> and)				
IT	Microorganism (Campylobacter pylori isolation and identification from, <b>urea</b> and acid in)				
IT	Stomach (Campylobacter pylori isolation from <b>biopsy</b> of, <b>urea</b> and acids in)				
IT	Feces (Campylobacter pylori isolation from, <b>urea</b> and acids in)				
IT	<b>Indicators</b> (acid-base, in Campylobacter pylori enrichment and isolation with <b>urea</b> and acids)				
IT	<b>7664-41-7, Ammonia</b> , biological studies RL: FORM (Formation, nonpreparative) (formation of, in Campylobacter pylori enrichment and isolation)				
IT	<b>9002-13-5, Urease</b> RL: FORM (Formation, nonpreparative) (formation of, Campylobacter pylori enrichment and isolation in relation to)				
IT	<b>57-13-6, Urea</b> , biological studies RL: BIOL (Biological study) (Campylobacter pylori enrichment and isolation in presence of acids and)				
IT	<b>7664-41-7, Ammonia</b> , biological studies RL: FORM (Formation, nonpreparative) (formation of, in Campylobacter pylori enrichment and isolation)				
RN	7664-41-7 HCAPLUS				
CN	Ammonia (8CI, 9CI) (CA INDEX NAME)				

NH3

IT **9002-13-5, Urease**  
RL: FORM (Formation, nonpreparative)



(formation of, *Campylobacter pylori* enrichment and isolation in relation to)

RN 9002-13-5 HCAPLUS

CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*


IT 57-13-6, Urea, biological studies

RL: BIOL (Biological study)

(*Campylobacter pylori* enrichment and isolation in presence of acids and)

RN 57-13-6 HCAPLUS

CN Urea (8CI, 9CI) (CA INDEX NAME)



L69 ANSWER 25 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 1987:433197 HCAPLUS

DN 107:33197

TI Bismuth composition for treatment of infectious **gastrointestinal** disorders

IN **Marshall, Barry James**

PA Australia

SO Eur. Pat. Appl., 11 pp.

CODEN: EPXXDW

DT Patent

LA English

IC ICM A61K031-60

ICS A61K033-00

CC 1-9 (Pharmacology)

Section cross-reference(s): 10

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 206627	A2	19861230	EP 1986-304409	19860610
	EP 206627	A3	19890405		
	EP 206627	B1	19920812		
	R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE				
	AT 79261	E	19920815	AT 1986-304409	19860610
	DE 3619733	A1	19870212	DE 1986-3619733	19860612
	DE 3619734	A1	19870212	DE 1986-3619734	19860612
	BE 904922	A1	19861215	BE 1986-216783	19860613
	JP 62048624	A2	19870303	JP 1986-138038	19860613
	JP 07094391	B4	19951011		
	US 5601848	A	19970211	US 1987-70857	19870708
PRAI	US 1985-744842	A	19850613		
	EP 1986-304409	A	19860610		

AB Treatment of title disorders in humans or lower animals comprises (1) testing for the presence of pyloric campylobacter or similar organism in the stomach; (2) on obtaining a pos. result, administering 50-5000 mg of Bi per day for 3-56 days, or until a neg. result is obtained on a diagnostic test. A preferred test for such infection is through detection of **urease** enzyme in the stomach. A human subject, suffering from peptic ulcer disease, was treated with 700 mg Bi (as Bi subsalicylate) per day for 35 days. The treatment resulted in healing of the peptic-ulcer crater.

ST infectious **gastrointestinal** disorder treatment bismuth

IT Campylobacter

Campylobacter pyloridis

(gastrointestinal infection by, bismuth treatment of)

IT Stomach, disease or disorder  
(atrophic gastritis, campylobacter detection in and bismuth treatment of)

IT Digestive tract  
(disease, infectious, bismuth treatment of)

IT Stomach, disease or disorder  
(mucosa, campylobacter detection in, bismuth treatment of)

IT Ulcer  
(peptic, infectious, bismuth treatment of)

IT 9002-13-5, Urease  
RL: ANT (Analyte); ANST (Analytical study)  
(detection of, in infectious gastrointestinal disorder, bismuth treatment in relation to)

IT 813-93-4, Bismuth citrate 1304-85-4 6591-56-6, Bismuth tartrate 7440-69-9, biological studies 14882-18-9, Bismuth subsalicylate 71156-53-1  
RL: BIOL (Biological study)  
(infectious gastrointestinal disorder treatment with)

IT 9002-13-5, Urease  
RL: ANT (Analyte); ANST (Analytical study)  
(detection of, in infectious gastrointestinal disorder, bismuth treatment in relation to)

RN 9002-13-5 HCAPLUS

CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 26 OF 34 HCAPLUS COPYRIGHT 2003 ACS

AN 1987:115736 HCAPLUS

DN 106:115736

TI Compositions, methods, and device for the detection of urease for the diagnosis of a Campylobacter pyloridis infection

IN Marshall, Barry James

PA Australia

SO Eur. Pat. Appl., 25 pp.  
CODEN: EPXXDW

DT Patent

LA English

IC ICM C12Q001-58

ICA G01N033-52; C12Q001-04

CC 7-1 (Enzymes)

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 204438	A2	19861210	EP 1986-303493	19860508
	EP 204438	A3	19870527		
	EP 204438	B1	19910306		
	R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE				
	AU 8657398	A1	19861120	AU 1986-57398	19850517
	AU 601363	B2	19900913		
	US 4748113	A	19880531	US 1985-744840	19850613
	CA 1274757	A1	19901002	CA 1986-508415	19860505
	AT 61414	E	19910315	AT 1986-303493	19860508
	ZA 8603605	A	19880127	ZA 1986-3605	19860515
	DK 8602283	A	19861118	DK 1986-2283	19860516
	DK 173710	B1	20010709		
	NO 8601966	A	19861118	NO 1986-1966	19860516
	NO 170091	B	19920601		
	NO 170091	C	19920909		
	BR 8602243	A	19870113	BR 1986-2243	19860516
	JP 62026000	A2	19870203	JP 1986-112427	19860516
	JP 06095960	B4	19941130		

PRAI AU 1985-611 A 19850517  
 US 1985-744840 A 19850613  
 EP 1986-303493 A 19860508

AB A reagent compn. for the detection of preformed **urease** for diagnosis of **gastrointestinal** disorders caused by *C. pyloridis* infection in a human or lower animal contains (1) **urea**, (2) a bactericide, (3) a **pH indicator** for detecting an increase in **pH**, and (4) water, where the compn. has a **pH** of .gtoreq.5.0, which is .gtoreq.1 unit lower than the **pKa** of the **indicator**. A reagent compn. contained **urea** 20, NaN3 1, **agar** 20 g/L, and **phenol red** 60 mg/L; the **pH** was adjusted to 5.50. Injection of a **sample** of vomitus from a human infant suspected of having **gastritis** into the **gelled** reagent compn. resulted in a change in the **color** of the **pH indicator** within 20 min.

ST reagent Campylobacter infection diagnosis; **urease** detection  
 Campylobacter infection diagnosis; **gastrointestinal** disorder  
 Campylobacter **urease** detection

IT Bactericides, Disinfectants, and Antiseptics  
 (in Campylobacter **urease** detection for  
**gastrointestinal** disorder diagnosis)

IT Campylobacter pyloridis  
 (**urease** of, detection of, in **gastrointestinal**  
 disorder diagnosis, reagents for)

IT **Indicators**  
 (acid-base, in Campylobacter **urease** detection for  
**gastrointestinal** disorder diagnosis)

IT Digestive tract  
 (disease, diagnosis of, by Campylobacter **urease** detection,  
 reagents for)

IT 26628-22-8, Sodium azide 29468-36-8  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (as bactericide, in reagent for Campylobacter **urease**  
 detection for **gastrointestinal** disorder diagnosis)

IT 143-74-8, **Phenol red**  
 RL: BIOL (Biological study)  
 (as **pH indicator**, in Campylobacter **urease**  
 detection for **gastrointestinal** disorder diagnosis)


IT 9002-13-5, **Urease**  
 RL: ANT (Analyte); ANST (Analytical study)  
 (detection of, of Campylobacter in **gastrointestinal** disorder  
 diagnosis, reagents for)

IT 57-13-6, **Urea**, uses and miscellaneous  
 RL: USES (Uses)  
 (in Campylobacter **urease** detection for  
**gastrointestinal** disorder diagnosis)

IT 143-74-8, **Phenol red**  
 RL: BIOL (Biological study)  
 (as **pH indicator**, in Campylobacter **urease**  
 detection for **gastrointestinal** disorder diagnosis)

RN 143-74-8 HCAPLUS


CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis- (9CI) (CA  
 INDEX NAME)



IT 9002-13-5, Urease  
 RL: ANT (Analyte); ANST (Analytical study)  
 (detection of, of Campylobacter in gastrointestinal disorder  
 diagnosis, reagents for)  
 RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

IT 57-13-6, Urea, uses and miscellaneous  
 RL: USES (Uses)  
 (in Campylobacter urease detection for  
 gastrointestinal disorder diagnosis)  
 RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)




L69 ANSWER 27 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
 AN 1982:139455 HCAPLUS  
 DN 96:139455  
 TI Dry nutritive media for the identification of Corynebacterium diphtheriae  
 AU Kovaleva, V. I.; Dzhililova, R. S.; Shtanchaeva, S. M.  
 CS Dagestan. Inst. Proizvod. Pitatel. Sred., Makhachkala, USSR  
 SO Razrab. Stand. Bakteriolog. Pitatel'nykh Sred (1980), 69-71. Editor(s):  
 Semenov, B. F.; Raskin, B. M. Publisher: Mosk. Nauchno-Issled. Inst.  
 Vaktsin Syvorotok im. I. I. Mechnikova, Moscow, USSR.  
 CODEN: 47IXA8  
 DT Conference  
 LA Russian  
 CC 10-2 (Microbial Biochemistry)  
 AB Dehydrated nutrient media were prepd. for cultivation and identification  
 of C. diphtheriae, C. pseudodiphtheriticum, and saprophytic diphtheroids.  
 Optimum medium for cultivation was composed of enzymic casein hydrolyzate  
 3, glucose 0.2, sucrose 1.0, NaCl 0.5, and acid fuchsin 0.009%, pH  
 7.6. Another culture medium, developed to test for urease  
 prodn. in Corynebacterium species, was composed of casein hydrolyzate 3.0,  
 glucose 1.0, NaCl 0.5, phenol red 0.0052, and  
 urea 1.0%, pH 7.1. These media were effective for  
 isolation, growth, and differentiation of Corynebacterium species.  
 ST Corynebacterium cultivation identification culture medium; urease  
 detn Corynebacterium culture medium  
 IT Caseins, compounds  
 RL: BIOL (Biological study)

DN 87:180311  
 TI Reagent mixture for determination of **urea**  
 IN Chang, Michael  
 PA USA  
 SO Ger. Offen., 20 pp.  
 CODEN: GWXXBX  
 DT Patent  
 LA German  
 IC G01N033-16  
 CC 9-6 (Biochemical Methods)  
 FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 2600146	A1	19770714	DE 1976-2600146	19760105
PRAI	DE 1976-2600146		19760105		

AB A reagent mixt. for the enzymic detn. of **urea** in biol. fluids is described that contains **urease**, an **indicator dye**, possibly a stabilizer, and a **buffer** whose pH rises with an increase in temp., and another **buffer** whose pH decreases with an increase in temp. Examples of the former type of **buffer** are the pyrophosphate derivs. and of the latter, substituted amines or phenols. Thus, a soln. contg. 10 mM triethanolamine, 10 mM pyrophosphate, 10 mM EDTA, 200 mM NaCl, and 0.3 mM **phenolsulfonephthalein** and distilled H<sub>2</sub>O was adjusted to pH 6-8. **Urease** was dissolved in some of the **buffer**-dye mixt. at an activity of .apprx.100 I.U./ml at 25.degree.. The change in absorbance at 560 nm was detd. for cuvetts contg. 2.8 mL of the enzyme-**buffer** and 0.1 mL of std. contg. 10, 50, or 100 mg **urea**-N/100 mL or 0.1 mL of serum or urine, and 0.1 mL of the **urease** soln. The reaction was allowed to go to completion, .gtoreq.5 h. The std. curve was linear up to 200 mg% **urea**-N.

ST serum urine **urea** detn; enzymic detn **urea**  
 IT **Buffer** substances and systems  
 (temp.-sensitive, for **urea** spectrometric detn.)  
 IT 57-13-6, analysis  
 RL: ANT (Analyte); ANST (Analytical study)  
 (detn. of, enzymic, temp. sensitive **buffer** reagent for)  
 IT 143-74-8 9002-13-5  
 RL: ANST (Analytical study)  
 (in **urea** detn.)  
 IT 57-13-6, analysis  
 RL: ANT (Analyte); ANST (Analytical study)  
 (detn. of, enzymic, temp. sensitive **buffer** reagent for)  
 RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)



IT 143-74-8 9002-13-5  
 RL: ANST (Analytical study)  
 (in **urea** detn.)  
 RN 143-74-8 HCAPLUS  
 CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis- (9CI) (CA INDEX NAME)



RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 30 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
 AN 1975:510869 HCAPLUS  
 DN 83:110869  
 TI Apparatus and method for analysis of urea  
 IN Gray, Don Norman; Keyes, Melvin H.; Semersky, Frank E.  
 PA Owens-Illinois, Inc., USA  
 SO Ger. Offen., 41 pp.  
 CODEN: GWXXBX  
 DT Patent  
 LA German  
 IC G01N  
 CC 9-6 (Biochemical Methods)  
 FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 2455970	A1	19750703	DE 1974-2455970	19741127
	DE 2455970	B2	19770721		
	US 3926734	A	19751216	US 1973-427322	19731221
	FR 2255602	A1	19750718	FR 1974-38065	19741120
	FR 2255602	B1	19780616		
	NL 7415252	A	19750624	NL 1974-15252	19741122
	NL 176308	B	19841016		
	NL 176308	C	19850318		
	CA 1039163	A1	19780926	CA 1974-214867	19741128
	GB 1494490	A	19771207	GB 1974-54031	19741213
	JP 50098396	A2	19750805	JP 1974-144361	19741216
	AU 7476761	A1	19760624	AU 1974-76761	19741223
PRAI	US 1973-427322		19731221		

AB Urea in aq. soln. may be detd. by hydrolyzing it to  $\text{NH}_4$   
 + by immobilized urease, treating the  $\text{NH}_4^+$  with  
 alkali, and passing the evolved  $\text{NH}_3$  through a hydrophobic  
 membrane where it is measured by a pH-sensitive electrode.  
 Thus, urease was immobilized on agar with CNBr. A  
 blood sample contg. urea was injected into a stream of  
 buffer that carried it into a chamber contg. the immobilized  
 enzyme. From there the sample, now contg.  $\text{NH}_4^+$  was  
 passed into another chamber where it was mixed with NaOH soln. and brought  
 into contact with a polypropylene film. The  $\text{NH}_3$  passing through  
 was measured by a Ag-AgCl electrode.

ST urea detn blood urine; enzymic detn urea; electrode  
 detn urea

IT Blood analysis  
 Urine analysis

(urea detn. in, enzymic app. for)  
 IT 57-13-6, analysis  
 RL: ANT (Analyte); ANST (Analytical study)  
 (detn. of, in biol. fluids, enzymic app. for)  
 IT 9002-13-5  
 RL: ANST (Analytical study)  
 (immobilized, for urea detn. in biol. fluids)  
 IT 57-13-6, analysis  
 RL: ANT (Analyte); ANST (Analytical study)  
 (detn. of, in biol. fluids, enzymic app. for)  
 RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)



IT 9002-13-5  
 RL: ANST (Analytical study)  
 (immobilized, for urea detn. in biol. fluids)  
 RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 31 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
 AN 1973:94524 HCAPLUS  
 DN 78:94524  
 TI Differential methods for identification of T-mycoplasmas based on demonstration of **urease**  
 AU Shepard, Maurice C.  
 CS Bacteriol. Div., Nav. Med. Field Res. Lab., Camp Lejeune, NC, USA  
 SO Journal of Infectious Diseases (1973), 127(Suppl.), S22-S25  
 CODEN: JIDIAQ; ISSN: 0022-1899  
 DT Journal  
 LA English  
 CC 9-4 (Biochemical Methods)  
 Section cross-reference(s): 10  
 AB The **indicator** reagent is 0.8% MnCl<sub>2</sub> contg. 1% **urea**.  
 The test is carried out at room temp. A pos. reaction for **urease**, i.e. the presence of **NH<sub>3</sub>** in T-mycoplasma colonies, is extremely rapid (within 5-10 sec), and the test must be performed on colonies <48 hr old. MnCl<sub>2</sub> is oxidized to insol., nearly colloidal MnO<sub>2</sub>, which ppts. on the surface of the colony, producing a dark, golden-brown colony when viewed by transmitted light under low-power microscopy. If a concn. of MnCl<sub>2</sub> >0.8% is used, the sensitivity of the reaction is markedly increased, and broad zones of reaction extending outward from the T-mycoplasma colony will be obsd. A differential **agar** medium was developed, contg. 0.03% MnSO<sub>4</sub>.  
 ST **urease** mycoplasma detn  
 IT Mycoplasma  
 (T-strain, identification of, **urease** in)  
 IT 9002-13-5  
 RL: ANST (Analytical study)  
 (in mycoplasma T-strain detection)  
 IT 9002-13-5  
 RL: ANST (Analytical study)  
 (in mycoplasma T-strain detection)  
 RN 9002-13-5 HCAPLUS  
 CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*


L69 ANSWER 32 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
AN 1965:45180 HCAPLUS  
DN 62:45180  
OREF 62:8051c-d  
TI **Urease** by diffusion assay  
AU Blain, J. A.; Caskie, M.  
CS Univ. Strathclyde, Glasgow, UK  
SO Chemistry & Industry (London, United Kingdom) (1965), (1), 17-18  
CODEN: CHINAG; ISSN: 0009-3068  
DT Journal  
LA English  
CC 57 (Enzymes)  
AB The assay is based on the liberation of **NH3** from **urea** in **agar gel** contg. **indicator**, the diam. of the **colored** zone which results being related to the concn. of the enzyme. The assay is as follows: to 150 ml. of water is added 2.25 g. **agar** and after boiling and cooling to 60.degree., 6 g. **urea**, 12 ml. of 0.04% cresol red in EtOH, and 1 ml. of 0.7% Na diethyldithiocarbamate are stirred in. The **agar** is poured into 140-mm. petri dishes to a depth of approx. 10 mm. and allowed to cool for 30 min. Vertical cups are cut with a 6-mm. cork borer, 12 in each plate, filled with enzyme soln. and covered at once with a 24 mm. cover slip. Diams. of the **colored** zones are measured in 90 min.  
IT Blood  
(analysis, detn. of aspartic aminotransferase)  
IT **9002-13-5, Urease**  
(detn. of)  
IT **9002-13-5, Urease**  
(detn. of)  
RN 9002-13-5 HCAPLUS  
CN Urease (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

L69 ANSWER 33 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
AN 1947:2439 HCAPLUS  
DN 41:2439  
OREF 41:499a-b  
TI **Urea** decomposition as a means of differentiating Proteus and paracolon cultures from each other and from Salmonella and Shigella types  
AU Christensen, W. Blake  
CS 1631 Mardall Blvd., San Antonio, TX  
SO Journal of Bacteriology (1946), 52, 461-6  
CODEN: JOBAAY; ISSN: 0021-9193  
DT Journal  
LA Unavailable  
CC 11C (Biological Chemistry: Microbiology)  
AB The following medium is recommended for detg., **urease** activity in organisms which cannot use **NH3** as the sole source of N: bacto-peptone 1, NaCl 5, KH2PO4 2, **phenol red** 0.012, **agar** 20, glucose 1, **urea** 2.0 g., and distd. water 100 cc. The **urea** is sterilized separately as a 20% soln. and added aseptically to the other constituents. The small quantity of peptone and the presence of glucose serve to counteract any alky. produced by peptone decompn. This medium shows Proteus, paracolon Aerobacter, and paracolon intermediates to be **urease** pos., while paracolon Escherichia, Salmonella, and Shigella are neg.  
IT Proteus  
(differentiation from paracolon bacteria, Salmonella and Shigella)  
IT Shigella  
(differentiation from Proteus and paracolon bacteria)



IT Bacteria  
 (paracolon, differentiation from Proteus, Salmonella and Shigella)  
 IT Aerobacter  
 Escherichia coli  
 (urea decompn. by)  
 IT Salmonella  
 (urea decompn. by, in differentiation from Proteus and  
 paracolon bacteria)  
 IT 57-13-6, Urea  
 (decompn. of, in bacteria differentiation)  
 IT 57-13-6, Urea  
 (decompn. of, in bacteria differentiation)  
 RN 57-13-6 HCAPLUS  
 CN Urea (8CI, 9CI) (CA INDEX NAME)



L69 ANSWER 34 OF 34 HCAPLUS COPYRIGHT 2003 ACS  
 AN 1944:39252 HCAPLUS  
 DN 38:39252  
 OREF 38:5861h-i,5862a  
 TI Manometric, titrimetric and **colorimetric** methods for measurement  
 of **urease** activity  
 AU Van Slyke, Donald D.; Archibald, Reginald M.  
 SO Journal of Biological Chemistry (1944), 154, 623-42  
 CODEN: JBCHA3; ISSN: 0021-9258  
 DT Journal  
 LA Unavailable  
 CC 11B (Biological Chemistry: Methods and Apparatus)  
 AB The original Van Slyke and Cullen (C. A. 10, 2356) manometric and  
 titrimetric procedures for measuring **urease** activity have been  
 modified for application to the more active **urease** preps. now  
 available. In order to stabilize jack bean **urease** at high  
 dilns. and to counteract the inactivation by Hg (present in manometric  
 app.), the **urease** has been dissolved in egg albumin (5%), the  
 Van Slyke-Neill chamber rinsed with albumin soln. before each analysis and  
 a concn. of 1% albumin maintained in the reacting **urea-**  
**urease** mixt. Three procedures have been described, gasometric (I)  
 titrimetric (II) and **colorimetric** (III) resp. In I, the enzyme  
 activity is measured by the rate of CO<sub>2</sub> formation, in II, by the rate of  
 NH<sub>3</sub> formation and in III, by the time required for enough (  
 NH<sub>4</sub>)<sub>2</sub>CO<sub>3</sub> to form to raise the pH of a phosphate  
 buffer from 6.8 to 7.7 (**phenol red**  
**indicator**).

=> sel hit rn  
 E31 THROUGH E36 ASSIGNED

=> fil reg  
 FILE 'REGISTRY' ENTERED AT 09:54:32 ON 30 JUN 2003  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2003 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
 provided by InfoChem.

STRUCTURE FILE UPDATES: 27 JUN 2003 HIGHEST RN 539020-41-2  
DICTIONARY FILE UPDATES: 27 JUN 2003 HIGHEST RN 539020-41-2

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 6, 2003

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. See HELP  
PROPERTIES for more information. See STNote 27, Searching Properties  
in the CAS Registry File, for complete details:  
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=> s e31-e36

1 9002-13-5/BI  
(9002-13-5/RN)

1 57-13-6/BI  
(57-13-6/RN)

1 143-74-8/BI  
(143-74-8/RN)

1 7664-41-7/BI  
(7664-41-7/RN)

1 9002-18-0/BI  
(9002-18-0/RN)

1 34487-61-1/BI  
(34487-61-1/RN)

L70 6 (9002-13-5/BI OR 57-13-6/BI OR 143-74-8/BI OR 7664-41-7/BI OR  
9002-18-0/BI OR 34487-61-1/BI)

=> d ide can tot

L71 ANSWER 1 OF 6 REGISTRY COPYRIGHT 2003 ACS

RN 34487-61-1 REGISTRY

CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis-, monosodium  
salt (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 3H-2,1-Benzoxathiole, phenol deriv.

CN Phenol, 4,4'-(3H-2,1-benzoxathiol-3-ylidene)bis-, S,S-dioxide, monosodium  
salt

OTHER NAMES:

CN Phenol red sodium

CN Phenol red, sodium salt

CN Phenolsulfonephthalein sodium

AR 27664-79-5

DR 115481-77-1, 27664-79-5

MF C19 H14 O5 S . Na


LC STN Files: BEILSTEIN\*, CA, CAPLUS, CASREACT, CHEMCATS, CHEMLIST, CSCHEM,  
IFICDB, IFIPAT, IFIUDB, MSDS-OHS, TOXCENTER, USPAT2, USPATFULL

(\*File contains numerically searchable property data)

Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*

(\*\*Enter CHEMLIST File for up-to-date regulatory information)

CRN (143-74-8)



● Na

21 REFERENCES IN FILE CA (1957 TO DATE)  
21 REFERENCES IN FILE CAPLUS (1957 TO DATE)

REFERENCE 1: 138:276108  
REFERENCE 2: 137:14966  
REFERENCE 3: 136:265236  
REFERENCE 4: 135:319066  
REFERENCE 5: 135:189831  
REFERENCE 6: 133:261080  
REFERENCE 7: 131:157574  
REFERENCE 8: 130:71632  
REFERENCE 9: 129:341459  
REFERENCE 10: 127:210387

L71 ANSWER 2 OF 6 REGISTRY COPYRIGHT 2003 ACS

RN 9002-18-0 REGISTRY

CN Agar (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Agar-agar (8CI)

OTHER NAMES:

CN Agar Agar Flake

CN Agargel

CN Agarpectin, mixt. with agarose

CN Agarose, mixt. with agarpectin

CN AX 30

CN Bacto-agar

CN Bengal gelatin

CN Bengal isinglass

CN Casitone

CN Ceylon isinglass

CN Chinese isinglass

CN D 100

CN D 100 (polysaccharide)

CN Deltagar LTS

CN Difco Bacto agar

CN Digenea simplex mucilage  
CN E 406  
CN GAM medium  
CN Gelose  
CN Hygicult TPC  
CN Ina Agar M 8  
CN Inagel N 6  
CN Japan agar  
CN Japan isinglass  
CN Kantenmatsu  
CN Laylor Carang  
CN Luxara 1253  
CN Oxoid III  
CN Oxoid L 11  
CN Phytagar  
CN S 10  
CN S 10 (polysaccharide)  
CN S 100  
CN S 100 (polysaccharide)  
CN T 1  
CN UP 16  
CN UP 37  
DR 63241-81-6  
MF Unspecified  
CI PMS, COM, MAN  
PCT Manual registration, Polyother, Polyother only  
LC STN Files: AGRICOLA, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CABA,  
CANCERLIT, CAPLUS, CASREACT, CHEMCATS, CHEMLIST, CIN, CSCHEM, DDFU,  
DIOGENES, DRUGU, EMBASE, HSDB\*, IFICDB, IFIPAT, IFIUDB, IPA, MEDLINE,  
MRCK\*, MSDS-OHS, NAPRALERT, NIOSHTIC, PIRA, PROMT, RTECS\*, SPECINFO,  
TOXCENTER, TULSA, USPAT2, USPATFULL  
(\*File contains numerically searchable property data)  
Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*  
(\*\*Enter CHEMLIST File for up-to-date regulatory information)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

5064 REFERENCES IN FILE CA (1957 TO DATE)  
88 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
5067 REFERENCES IN FILE CAPLUS (1957 TO DATE)

REFERENCE 1: 139:12314  
REFERENCE 2: 139:11465  
REFERENCE 3: 139:7823  
REFERENCE 4: 139:6171  
REFERENCE 5: 139:6069  
REFERENCE 6: 139:6042  
REFERENCE 7: 139:5852  
REFERENCE 8: 138:406893  
REFERENCE 9: 138:406789  
REFERENCE 10: 138:406227

L71 ANSWER 3 OF 6 REGISTRY COPYRIGHT 2003 ACS  
RN 9002-13-5 REGISTRY  
CN Urease (8CI, 9CI) (CA INDEX NAME)

## OTHER NAMES:

CN E.C. 3.5.1.5  
CN Urea amidohydrolase  
CN **Urease LF**  
MF Unspecified  
CI COM, MAN  
LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BIOBUSINESS, BIOSIS, BIOTECHNO,  
CA, CABA, CAPLUS, CASREACT, CBNB, CEN, CHEMCATS, CHEMLIST, CIN, CSCHEM,  
CSNB, DDFU, DRUGU, EMBASE, IFICDB, IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*,  
MSDS-OHS, NAPRALERT, PIRA, PROMT, RTECS\*, TOXCENTER, TULSA, USPAT2,  
USPATFULL, VTB  
(\*File contains numerically searchable property data)  
Other Sources: EINECS\*\*, TSCA\*\*  
(\*Enter CHEMLIST File for up-to-date regulatory information)

## \*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

6903 REFERENCES IN FILE CA (1957 TO DATE)  
211 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
6909 REFERENCES IN FILE CAPLUS (1957 TO DATE)

REFERENCE 1: 139:12811  
REFERENCE 2: 139:6211  
REFERENCE 3: 139:6204  
REFERENCE 4: 139:6009  
REFERENCE 5: 139:4736  
REFERENCE 6: 139:3193  
REFERENCE 7: 139:3149  
REFERENCE 8: 138:401056  
REFERENCE 9: 138:398525  
REFERENCE 10: 138:398447

L71 ANSWER 4 OF 6 REGISTRY COPYRIGHT 2003 ACS

RN 7664-41-7 REGISTRY

CN **Ammonia (8CI, 9CI)** (CA INDEX NAME)

## OTHER NAMES:

CN Ammonia gas  
CN Ammonia-14N  
CN Nitro-Sil  
CN R 717  
CN Refrigerent R717  
CN Spirit of Hartshorn  
FS 3D CONCORD  
DR 8007-57-6, 208990-07-2, 214478-05-4  
MF H3 N  
CI COM  
LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, AQUIRE, BIOBUSINESS, BIOSIS,  
BIOTECHNO, CA, CABA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB, CEN,  
CHEMCATS, CHEMINFORMRX, CHEMLIST, CHEMSAFE, CIN, CSCHEM, CSNB, DDFU,  
DETERM\*, DIOGENES, DIPPR\*, DRUGU, EMBASE, ENCOMPLIT, ENCOMPLIT2,  
ENCOMPPAT, ENCOMPPAT2, GMELIN\*, HSDB\*, IFICDB, IFIPAT, IFIUDB, IPA,  
MEDLINE, MRCK\*, MSDS-OHS, NIOSHTIC, PDLCOM\*, PHARMASEARCH, PIRA, PROMT,  
RTECS\*, SPECINFO, TOXCENTER, TULSA, ULIDAT, USAN, USPAT2, USPATFULL,  
VETU, VTB  
(\*File contains numerically searchable property data)

Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*  
(\*\*Enter CHEMLIST File for up-to-date regulatory information)

NH3

114987 REFERENCES IN FILE CA (1957 TO DATE)  
1616 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
115045 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
1 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 139:16844

REFERENCE 2: 139:16839

REFERENCE 3: 139:16758

REFERENCE 4: 139:16757

REFERENCE 5: 139:16658

REFERENCE 6: 139:16647

REFERENCE 7: 139:16633

REFERENCE 8: 139:16073

REFERENCE 9: 139:16072

REFERENCE 10: 139:16054

L71 ANSWER 5 OF 6 REGISTRY COPYRIGHT 2003 ACS

RN 143-74-8 REGISTRY

CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis- (9CI) (CA  
INDEX NAME)

OTHER CA INDEX NAMES:

CN 3H-2,1-Benzoxathiole, phenol deriv.

CN Phenol, 4,4'-(3H-2,1-benzoxathiol-3-ylidene)bis-, S,S-dioxide

CN Phenol, 4,4'-(3H-2,1-benzoxathiol-3-ylidene)di-, S,S-dioxide (8CI)

OTHER NAMES:

CN .alpha.-Hydroxy-.alpha.,.alpha.-bis(p-hydroxyphenyl)-o-toluenesulfonic  
acid .gamma.-sultone

CN 3,3-Bis(p-hydroxyphenyl)-2,1,3H-benzoxathiole 1,1-dioxide

CN 3H-2,1-Benzoxathiole, 3,3-bis(4-hydroxyphenyl)-, 1,1-dioxide

CN Fenolipuna

CN Phenol red

CN Phenolsulfonephthalein

CN Phenolsulfonphthalein

CN Phenolsulphonphthalein

CN PSP

CN PSP (indicator)

CN Sulfonphthal

CN Sulphental

CN Sulphonthal

CN TF-R 2

FS 3D CONCORD


DR 2877-88-5

MF C19 H14 O5 S

CI COM

LC STN Files: AGRICOLA, ANABSTR, AQUIRE, BEILSTEIN\*, BIOBUSINESS, BIOSIS,  
BIOTECHNO, CA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CEN, CHEMCATS,  
CHEMLIST, CIN, CSCHEM, DDFU, DIOGENES, DRUGU, EMBASE, HODOC\*, IFICDB,

IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*, MSDS-OHS, NIOSHTIC, PIRA, PROMT,  
RTECS\*, SPECINFO, TOXCENTER, ULIDAT, USAN, USPAT2, USPATFULL  
(\*File contains numerically searchable property data)  
Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*  
(\*\*Enter CHEMLIST File for up-to-date regulatory information)



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1262 REFERENCES IN FILE CA (1957 TO DATE)  
31 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
1265 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
27 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 139:8461  
REFERENCE 2: 138:390732  
REFERENCE 3: 138:363805  
REFERENCE 4: 138:358106  
REFERENCE 5: 138:353859  
REFERENCE 6: 138:353073  
REFERENCE 7: 138:352809  
REFERENCE 8: 138:351453  
REFERENCE 9: 138:314635  
REFERENCE 10: 138:309402

L71 ANSWER 6 OF 6 REGISTRY COPYRIGHT 2003 ACS

RN 57-13-6 REGISTRY

CN Urea (8CI, 9CI) (CA INDEX NAME)

OTHER NAMES:

CN Aquacare  
CN Aquadrate  
CN B-I-K  
CN Basodexan  
CN Benural 70  
CN Carbamide  
CN Carbamimidic acid  
CN Carbonyl diamide  
CN Elaqua XX  
CN Eucerin 10% Urea Lotion


CN Hyanit  
 CN Isourea  
 CN Keratinamin  
 CN Keratinamin Kowa  
 CN Nutraplus  
 CN Onychomal  
 CN Optigen 1200  
 CN Pastaron  
 CN Pastaron 10  
 CN Pastaron 20  
 CN Pastaron 20 soft  
 CN Pseudourea  
 CN UR  
 CN Urea perhydrate  
 CN Ureaphil  
 CN Ureophil  
 CN Urepeal  
 CN Urepeal L  
 CN Urepearl  
 CN Urevert  
 CN Varioform II  
 FS 3D CONCORD  
 DR 30535-50-3  
 MF C H4 N2 O  
 CI COM

LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, AQUIRE, BEILSTEIN\*, BIOBUSINESS,  
 BIOSIS, BIOTECHNO, CA, CABA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB,  
 CEN, CHEMCATS, CHEMINFORMRX, CHEMLIST, CIN, CSCHEM, CSNB, DDFU,  
 DETHERM\*, DIOGENES, DIPPR\*, DRUGU, EMBASE, ENCOMPLIT, ENCOMPLIT2,  
 ENCOMPPAT, ENCOMPPAT2, GMELIN\*, HODOC\*, HSDB\*, IFICDB, IFIPAT, IFIUDB,  
 IPA, MEDLINE, MRCK\*, MSDS-OHS, NAPRALERT, NIOSHTIC, PDLCOM\*, PHAR, PIRA,  
 PROMT, RTECS\*, SPECINFO, SYNTHLINE, TOXCENTER, TULSA, ULIDAT, USAN,  
 USPAT2, USPATFULL, VETU, VTB

(\*File contains numerically searchable property data)

Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*

(\*\*Enter CHEMLIST File for up-to-date regulatory information)



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

63891 REFERENCES IN FILE CA (1957 TO DATE)  
 3044 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
 63935 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
 9 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 139:16820

REFERENCE 2: 139:15273

REFERENCE 3: 139:13047

REFERENCE 4: 139:12811

REFERENCE 5: 139:12682

REFERENCE 6: 139:12670



REFERENCE 7: 139:12298

REFERENCE 8: 139:12210

REFERENCE 9: 139:12084

REFERENCE 10: 139:11970

=> fil wpiX

FILE 'WPIX' ENTERED AT 10:27:58 ON 30 JUN 2003

COPYRIGHT (C) 2003 THOMSON DERWENT

FILE LAST UPDATED: 24 JUN 2003 <20030624/UP>  
MOST RECENT DERWENT UPDATE: 200340 <200340/DW>  
DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> NEW WEEKLY SDI FREQUENCY AVAILABLE --> see NEWS <<<

>>> SLART (Simultaneous Left and Right Truncation) is now  
available in the /ABEX field. An additional search field  
/BIX is also provided which comprises both /BI and /ABEX <<<

>>> PATENT IMAGES AVAILABLE FOR PRINT AND DISPLAY <<<

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES,  
SEE <http://www.derwent.com/dwpi/updates/dwpicov/index.html> <<<

>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,  
PLEASE VISIT:  
[http://www.stn-international.de/training\\_center/patents/stn\\_guide.pdf](http://www.stn-international.de/training_center/patents/stn_guide.pdf) <<<

>>> FOR INFORMATION ON ALL DERWENT WORLD PATENTS INDEX USER  
GUIDES, PLEASE VISIT:  
[http://www.derwent.com/userguides/dwpi\\_guide.html](http://www.derwent.com/userguides/dwpi_guide.html) <<<

=> d all abeq tech abex tot

L113 ANSWER 1 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 2003-120556 [11] WPIX

DNN N2003-096041 DNC C2003-031141

TI Positive response biosensor for detecting analyte in environment, has  
first reaction system having enzyme and substrate for enzyme, and second  
reaction system that produces detectable state when enzyme is inhibited.

DC B04 D16 S03

IN ERBELDINGER, M; LEJEUNE, K E

PA (ERBE-I) ERBELDINGER M; (LEJE-I) LEJEUNE K E; (AGEN-N) AGENTASE LLC

CYC 100

PI WO 2002090577 A2 20021114 (200311)\* EN 29p C12Q001-34

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TR TZ UG ZM ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK  
DM DZ EC EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR  
KZ LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ OM PH PL PT  
RO RU SD SE SG SI SK SL TJ TM TN TR TT TZ UA UG UZ VN YU ZA ZM ZW

US 2002182662 A1 20021205 (200311) C12Q001-34

ADT WO 2002090577 A2 WO 2002-US13692 20020502; US 2002182662 A1 US 2001-850686  
20010507

PRAI US 2001-850686 20010507

IC ICM C12Q001-34

ICS C12M001-00; C12M001-34; C12Q001-00; C12Q001-44; G01N033-53

AB WO 200290577 A UPAB: 20030214

NOVELTY - Sensor for detecting analyte (A) comprises first reaction system

(RS1) with first enzyme (E1) and substrate for E1, where (A) inhibits E1, and second reaction system (RS2) that produces first detectable state (DS1) when E1 is inhibited, or RS1 with E1 or first substrate (S1), where (A) is substrate for E1 if ES1 includes E1/S1, and RS2 that produces DS1 when (A) is below certain concentration.

DETAILED DESCRIPTION - A sensor (I) for detecting an analyte in an environment, comprises:

(a) a first reaction system including a first enzyme and a substrate for the first enzyme, where the analyte inhibits the first enzyme, and a second reaction system that reacts to produce a first detectable state when the first enzyme is inhibited; or

(b) a first reaction system including a first enzyme or a first substrate, where the analyte is a substrate for the first enzyme if the first reaction system includes the first enzyme or the first substrate, and a second reaction system that reacts to produce a first detectable state when the analyte is below a certain concentration.

USE - (I) is useful for detecting an analyte such as a nerve agent in an environment (claimed). (I) is useful for detecting the presence of an enzyme inhibitor or a substrate deficiency with a positive signal in form of, for example, changing pH or color, or for monitoring the absence of an enzymatic reaction as a result of inhibitor presence or substrate deficiency.

ADVANTAGE - Compared to the prior art detection of nerve agents where detection relies on negative response of the inhibition of the cholinesterase enzyme, (I) provides a positive response signal which provides a changing signal in the presence of contamination. For example, in the case of inhibitor detection or the detection of compound/substrate deficiency, (I) improves the prior art by providing a positive signal even in the absence of an enzymatic reaction. Prior art sensors for detection of nerve agents include cholinesterase paired with its respective substrate. When nerve agents are present, cholinesterase is inhibited and the signal is retarded or nonexistent. Only in the absence of nerve agents does the enzymatically catalyzed reaction of the substrate occur. In (I), a second enzyme such as **urease** is added to a butyryl cholinesterase-based sensor. Hydroxide ions resulting from the formation of ammonium during hydrolysis of **urea** neutralize the protons produced during the hydrolysis of cholinesterase substrate (butyrylcholine). When nerve gas agents are absent both enzymatic systems are active and no pH change occurs. When an agent is present, hydroxide ions resulting from the hydrolysis of **urea** are not neutralized because butyryl cholinesterase is inhibited. The pH of the sensor then rises, resulting in positive signal.

Dwg.0/9

FS CPI EPI

FA AB; DCN

MC CPI: B04-L05; **B11-C07B1**; B11-C08E3; B12-K04E; D05-A01A2;  
D05-A01B3; D05-H09

EPI: S03-E03X; S03-E04E; **S03-E14H4**

TECH UPTX: 20030214

TECHNOLOGY FOCUS - BIOTECHNOLOGY - Preferred Sensor: In (I), the reaction of the first reaction system produces a second detectable state, different from the first detectable state. The reaction of the first reaction system causes pH to change in a first direction and the reaction of the second reaction system causes pH to change in a second direction, opposite of the first direction. The second reaction system comprises a second enzyme and a substrate for the second enzyme. The first enzyme is a hydrolase or cholinesterase, and the second enzyme is a different hydrolase. The first detectable change is a colorimetric change. The reaction of the first reaction system produces a second detectable state, different from the first detectable state. The first detectable state arises from the presence of a first pH sensitive dye producing a colorimetric change and the second detectable state is a colorimetric change different from the colorimetric change of the first

detectable state. The reaction of the first reaction system causes a first colorimetric change and the reaction of the second reaction system causes a second colorimetric change, where the second colorimetric change is different from the first colorimetric change. The reaction of the first reaction system causes **pH** to change in a first direction and the reaction of the second reaction system causes a **pH** sensitive colorimetric change when the first enzyme is inhibited or when the analyte is below a certain concentration. The first and second enzyme is immobilized in a polymer medium. The reaction of the analyte catalyzed by the first enzyme produces a second detectable state, different from the first detectable state. The reaction of the analyte catalyzed by the first enzyme causes **pH** to change in a first direction and the reaction of the second reaction system causes **pH** to change in a second direction, opposite of the first direction. (I) comprises a first reaction system that is reduced in reactivity by the presence of the analyte, and at least a second reaction system that reacts to produce a first detectable state when the first reaction system is inhibited. (I) comprises a first reaction system including a first compound that produces a reaction with the analyte, and at least a second reaction system that reacts to produce a first detectable state when the analyte is below a certain concentration.

ABEX

UPTX: 20030214

EXAMPLE - Detection of diisopropyl fluorophosphate (DFP) using a positive response enzymatic biosensor with butyrylcholinesterase (BChE) immobilized in polyurethane, **urease** and a **pH**-sensitive dye (cresol red) was as follows: Hydroxide ions resulting from the formation of **ammonia** neutralized any protons produced during hydrolysis of butyrylcholine. No color change from the original yellow was observed as a result of stabilized **pH** when both enzymes were active. In the presence of DFP, however BChE was significantly inhibited while **urease** remained active. Only hydroxide ions were produced and **pH** increased accordingly. Increasing **pH** resulted in a color change of incorporated dye and the sensor changed from yellow to red. The color change was easily recognized by the naked eye. To remove any subjectivity from the experimental procedures, a solid-phase Minolta CM-500d solid spectrophotometer was used to monitor the sensor's color change. This unit used a three-dimensional color coordinate system to define colors and intensity. The biopolymer containing cresol red developed yellow color when **pH** was below 7.0 and turned to red at a **pH** around 8.8. Each kinetic reaction was performed in duplicate. It was clear that a positive response was observed in the presence of DFP, a powerful inhibitor of the cholinesterase sensing enzyme used in this sensor construct.

L113 ANSWER 2 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 2003-013562 [01] WPIX

DNN N2003-009786 DNC C2003-003115

TI Method for determining helicobacter pylori-associated intragastral **urease** activity.

DC B04 D16 S03

IN AKHMETSHIN, R Z; KHASANOV, R SH; LOGINOVSKAYA, V V; MELNIKOVA, Z M; NIZHEVICH, A A; SATAEV, V U

PA (NIZH-I) NIZHEVICH A A; (UYBA-R) UNIV BASHKIR MED

CYC 1

PI RU 2189591 C1 20020920 (200301)\* G01N033-48 &lt;--

ADT RU 2189591 C1 RU 2001-102196 20010124

PRAI RU 2001-102196 20010124

IC ICM G01N033-48

ICS G01N033-49

AB RU 2189591 C UPAB: 20030101

NOVELTY - Method for detecting the degree of bacterial seeding volume of gastric mucosa at Helicobacter gastritis, gastroduodenitis and ulcerous disease.

DETAILED DESCRIPTION - A **biopsy** fragment of gastric mucosa is taken and put into commercial solution. Incubation mixture is subjected for exposure, PEC- colorimetry is conducted at 540 nm wave length to compare optic density with incubation time of **biopsy** fragment and the weight of **biopsy** fragment (units of optic density/mg **biopsy** material/min). At **urease** activity values ranged 11.5-4 U one should detect a low degree of bacterial seeding volume of gastric mucosa, at its value within 5-10 U a moderate degree is detected and in case its value ranges 11-19 U a high degree of *Helicobacter pylori* seeding volume is concluded on. The method is of high specificity, enables user to conduct a semi-quantitative analysis of bacterial seeding volume of gastric mucosa.

USE - Medicine, medicinal microbiology and gastroenterology.

ADVANTAGE - Higher efficiency.

Dwg.0/0

FS CPI EPI

FA AB

MC CPI: B04-F10; B04-L01; B11-A01; B11-A02; B11-C08E1; B11-C08E3; B12-K04A4;  
B12-K04E; D05-A02; D05-H04; D05-H08; D05-H09  
EPI: S03-E14H; S03-E14H1

L113 ANSWER 3 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 2001-464387 [50] WPIX

CR 2001-335013 [35]

DNC C2001-140251

TI A device for the in vivo detection of **urease**-producing *Helicobacter* in the stomach.

DC B04 D16

IN MARSHALL, B

PA (MARS-I) MARSHALL B

CYC 1

PI US 2001012623 A1 20010809 (200150)\* 6p C12Q001-04 <--  
US 6479278 B2 20021112 (200278) C12M001-34

ADT US 2001012623 A1 Cont of US 1995-489816 19950613, CIP of US 1997-832332  
19970326, US 2001-824870 20010403; US 6479278 B2 Cont of US 1995-489816  
19950613, CIP of US 1997-832332 19970326, US 2001-824870 20010403

FDT US 2001012623 A1 CIP of US 6228605; US 6479278 B2 CIP of US 6228605

PRAI US 2001-824870 20010403; US 1995-489816 19950613; US 1997-832332  
19970326

IC ICM C12M001-34; C12Q001-04

AB US2001012623 A UPAB: 20021204

NOVELTY - A diagnostic device for the in vivo detection of **urease**-producing *Helicobacter* in the upper stomach, is new.

DETAILED DESCRIPTION - A diagnostic device for the detection of **urease** producing *Helicobacter* in a subjects stomach comprising a soluble carrier containing a combination of a **pH indicator** (pHI1) with a **pH** range of 5.5-9.0 (pHI1 has an first indicium to **indicate** an alkaline **pH** range and a second indicium to **indicate** an acidic **pH** range) and a second **pH indicator** (pHI2) with a **pH** range of 5.5-9.0 (pHI2 has a first indicium to **indicate** an acidic **pH** and a third indicium to **indicate** an alkaline **pH** range, and a reagent which reacted with **urease** to produce **ammonia**). The pHI1 first indicium and the pHI2 first indicium are the same. The pHI1 second indicium and the pHI2 third indicium are different from one another and from the pHI1 and pHI2 first indicia. The pHI1 and pHI2 **indicator** combination react to a presence or absence of **urease** producing *Helicobacter* by change, or lack of change of indicia.

If pHI1 and pHI2 combine to **indicate** an acidic **pH**, this **indicates** an absence of the *Helicobacter* (the stomach is acidic and there are no **urease**-producing *Helicobacter*. If the pHI1 and pHI2 combine to **indicate** an alkaline **pH**, this

**indicates** that the stomach is alkaline and no determination can be made, therefore producing a false positive result.

If the **pHI1 indicates** an acidic **pH** and the **pHI2 indicates** an alkaline **pH**, this **indicates** the presence of **ammonia** and the presence of **urease** producing **Helicobacter**.

USE - The device is used for the in vivo detection of **urease** -producing **Helicobacter** in the stomach.

ADVANTAGE - The device is used in vivo, eliminating the need for a **biopsy**.

Dwg.0/1

FS

CPI

FA

AB; DCN

MC

CPI: B04-C01; B04-F10; B04-L01; B04-N04; B11-A02; B11-C08E1; B11-C08E3;  
**B12-K04A4**; B12-K04E; D05-A02; D05-H04; D05-H08; D05-H09;  
D05-H10

TECH

UPTX: 20010905

TECHNOLOGY FOCUS - INSTRUMENTATION AND TESTING - Preferred Device: The device further comprises first and second dense carriers which are soluble in gastric fluids and have densities that cause them to descend through the stomach's fluids to the stomach's gastric mucosa. The first dense carrier is combined with the **pHI1** and the second dense carrier is combined with the **pHI2**. The container is a soluble capsule comprising the first carrier and second carrier. The dense carrier materials sorb the **indicators** and dissolve in the gastric fluids within 5 minutes after reaching the stomach's gastric mucosa. The dense carrier materials are in the form of beads which facilitate the dispersal of the **indicators** over the mucosa.

The indicium is color. The **pHI1** first indicium is one color at an acidic **pH** and the second indicium is a second color at an alkaline **pH**. The **pHI2** first indicium is one color at an acidic **pH** and the second indicium is a third color at an alkaline **pH**. each of the **pHI1** first indicium, and the **pHI2** first indicium can be the same color and/or the **pHI1** second indicium and the **pHI2** third indicium are different colors from one another and from the **pHI1** first indicium and the **pHI2** first indicium.

The reagent is **urea**.

ABEX

UPTX: 20010905

ADMINISTRATION - The device may be swallowed by the patient.

EXAMPLE - Beads comprising bromothymol blue **indicator**, buffer ( **pH** 6) and sugar beads and **phenol red**

**indicator**, buffer (**pH** 6), sugar beads and **urea**

were encapsulated into a quick dissolving gelatin capsule for delivery into the stomach in mass and undiluted.

L113 ANSWER 4 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 2001-354606 [37] WPIX

DNN N2001-257673 DNC C2001-109767

TI Gastrointestinal **sampling** device for diagnosis of certain gastrointestinal pathogens, comprises drag material for obtaining gastrointestinal **sample**, and protective sheath for deployment about the drag material.

DC A96 P31

IN **MARSHALL, B**; WONG, A M

PA (UYWA-N) UNIV WESTERN AUSTRALIA

CYC 94

PI WO 2001015604 A1 20010308 (200137)\* EN 51p A61B010-00

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TZ UG ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM  
DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC  
LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE  
SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2000068115 A 20010326 (200137) A61B010-00  
 JP 2003508106 W 20030304 (200319) 43p A61B010-00  
 ADT WO 2001015604 A1 WO 2000-AU1047 20000831; AU 2000068115 A AU 2000-68115  
 20000831; JP 2003508106 W WO 2000-AU1047 20000831, JP 2001-519821 20000831  
 FDT AU 2000068115 A Based on WO 200115604; JP 2003508106 W Based on WO  
 200115604  
 PRAI AU 1999-4609 19991213; AU 1999-2541 19990831  
 IC ICM A61B010-00  
 AB WO 200115604 A UPAB: 20010704

NOVELTY - A gastrointestinal **sampling** device (10) comprises a  
 drag material (12) for obtaining a gastrointestinal **sample**; and  
 a protective sheath (22) for deployment about the drag material. The drag  
 material is enclosed by the protective sheath upon removal from the  
 gastrointestinal tract.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for:

(a) a method of gastrointestinal **sampling**, comprising  
 swallowing the gastrointestinal **sampling** device; allowing a time  
 for the drag material to obtain the gastrointestinal **sample**;  
 withdrawing the drag material so that upon withdrawal, the protective  
 sheath encloses the drag material; and recovering the gastrointestinal  
**sample** for testing; and

(b) a method of manufacturing a gastrointestinal **sampling**  
 device, comprising encasing the drag material and the protective sheath in  
 a capsule.

USE - The inventive device is used for the diagnosis of certain  
 gastrointestinal pathogens (claimed). It is useful for obtaining  
**samples** from under the gastric mucus and between the epithelial  
 cells of the stomach.

ADVANTAGE - The inventive device increases the epithelial cells  
 removed from the stomach lining without causing additional discomfort to  
 the patient. Further, it is capable of **sampling** microorganisms  
 from specific regions of the gastrointestinal tract without becoming  
 contaminated with microorganisms from other regions.

DESCRIPTION OF DRAWING(S) - The figure shows a front side view of the  
 gastrointestinal **sampling** device.

Gastrointestinal **sampling** device 10

Drag material 12

Capsule 14

Weight 16

Glue 18

Protective sheath 22

Dwg.1/11

FS CPI GMPI

FA AB; GI

MC CPI: A12-V03C2

TECH UPTX: 20010704

TECHNOLOGY FOCUS - INSTRUMENTATION AND TESTING - Preferred Component: The  
 drag material includes a **pH indicator** or

**urease indicator**, and is folded within a capsule. The  
 capsule (14) carries the drag material and the protective sheath. It is  
 constructed of a non-gelatin dissolvable material, and comprises at least  
 two parts joined together with water-soluble glue (18). The protective  
 sheath is deployed about the drag material by movement from a retracted  
 position to an extended position. It is assisted in deployment by the use  
 of a spring device. The retracted position is held in place by an edible  
 glue. The spring device is attached with adhesive to the inner surface of  
 the protective sheath. A non-absorbent filament has two ends. The first  
 end is attached to one end of the protective sheath. The second end is  
 attached to a weight (16), preferably a dissolvable weight. The weight  
 assists with the extension of the drag material in the stomach of a  
 patient. A rubber ring is attached to the sheath, and has a similar  
 diameter to the sheath and assists in the deployment of the sheath.  
 Preferred Properties: The capsule is more than 2 cm long, and is no more

than 0.9 cm in diameter.

TECHNOLOGY FOCUS - POLYMERS - Preferred Material: The drag material is an absorbent string, cotton, **sampling** cloth, wool, acrylic, nylon, plastic, chain links, and/or finely woven metal. It is 50% wool and 50% acrylic, and is coated with a bacterial adherent which is poly-L-lysine. The spring device is a nylon line or a Teflon coated stainless steel thread.

L113 ANSWER 5 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 2001-335013 [35] WPIX

CR 1995-178631 [23]; 2001-464387 [50]

DNC C2001-103414

TI Detecting **urease**-producing *Helicobacter* in a patient's stomach, by administering encapsulated dense carrier treated with reagent **indicators**, one containing **urea**, and observing color changes in the gastric mucosa.

DC B04 D16

IN MARSHALL, B J

PA (MARS-I) MARSHALL B J

CYC 1

PI US 6228605 B1 20010508 (200135)\* 8p C12Q001-04 <--

ADT US 6228605 B1 CIP of US 1993-142600 19931028, Cont of US 1995-489816 19950613, US 1997-832332 19970326

PRAI US 1995-489816 19950613; US 1993-142600 19931028; US 1997-832332 19970326

IC ICM C12Q001-04

AB US 6228605 B UPAB: 20010905

NOVELTY - Detecting **urease**-producing *Helicobacter* in a patient's stomach using a dense carrier (C) which is divided into 2 separate groups which are combined with separate reagent **indicators**, one of which contains **urea** (U), administering (C) and (U) encapsulated in a solid capsule (SC) to the patient, dissolving SC in stomach fluids, contacting the reagents with a gastric mucosa and observing color changes.

DETAILED DESCRIPTION - Detecting (M), in vivo, the presence or absence of **urease** producing *Helicobacter* in a patient's stomach involves:

(a) administering to a patient a pharmaceutically acceptable soluble container containing a combination comprising a first **indicator** having a pH indicium range of from about 5.5-9.0 and having a first indicium for **indicating** an acidic pH range and a second indicium for **indicating** an alkaline pH, and a second **indicator** combination, where the second **indicator** combination has a second pH **indicator** having a pH indicium range of from about 5.5-9.0 and having a second pH **indicator** first indicium for **indicating** an acidic pH range and a second pH **indicator** third indicium for **indicating** an alkaline pH range, and a reagent to react with **urease** in the stomach to form an alkaline product, the first pH **indicator** first indicium and the second pH **indicator** combination first indicium being the same, the first pH **indicator** second indicium and the second pH **indicator** combination third indicium being different from one another, from the first pH **indicator** first indicium and from the second pH **indicator** first indicium;

(b) dissolving the soluble container in the patients stomach fluids;

(c) contacting the patients gastric mucosa with the first pH **indicator** and the second **indicator** combination; and

(d) observing the first pH **indicator** and the second **indicator** combination in the patient's stomach where if:

(i) the first pH **indicator** first indicium and the second **indicator** combination first indicium **indicate** an acidic pH range, then the stomach is acidic,

indicating an absence of **urease** producing *Helicobacter*;

(ii) the first **pH indicator** second indicium and the second **indicator** combination third indicium **indicate** an alkaline **pH** range, then the stomach is alkaline, and thus no determination can be made regarding the presence or absence of **urease** producing *Helicobacter*; or

(iii) the first **pH indicator** first indicium **indicates** an acidic **pH** range and the second **indicator** combination third indicium **indicates** an alkaline **pH** range, then the stomach is acidic **indicating** the presence of **urease** producing *Helicobacter*.

USE - (M) Is useful for diagnosing gastrointestinal disorders caused by **urease** producing *Helicobacter* by determining the presence or absence of **urease** within a subject's stomach by:

(a) administering to the subject between approximately 1 and 20 g of **urea**/kg of dense, pharmaceutically acceptable carrier, the carrier having a density greater than body fluids, the **urea** being carried by the dense carrier;

(b) drinking a predetermined quantity of a liquid, delivering the capsule through stomach fluids to the subject's gastric mucosa, the dense carrier causing the first **pH indicator**, the second **pH indicator** and the **urea** to descent through the stomach fluids;

(c) dissolving the capsule in gastric juices contained in the subjects stomach, thus placing the carrier, the **pH indicators** and the **urea** in direct contact with the gastric mucosa;

(d) reacting the **urea** with any **urease** present to produce **ammonia**, thus raising the **pH** proximate to the **indicators** within the subject's stomach; and

(e) viewing the first **pH indicator** indicium and the second **pH indicator** indicium for an **indication** of **pH** change, the **pH** change **indicating** the absence or presence of *Helicobacter*, where when viewed if:

(i) the first indicium of the first **pH indicator** and the first indicium of the second **pH indicator** are a color that **indicate** an acidic range, then there is an absence of **urease** and a negative **indication** of the presence of the *Helicobacter*;

(ii) the second indicium of the first **pH indicator** and the third indicium of the second **pH indicator** are a color which **indicate** an alkaline **pH** range, then no determination regarding a gastrointestinal disorder can be made; or

(iii) the first indicium of the first **indicator** is a color that **indicates** an acidic range and the third indicium of the second **pH indicator** is a color that **indicates** **urea** in the second **pH indicator** combination is reacting with the **urease** to create an alkaline **pH**, then there is a positive **indication** of a presence of *Helicobacter*, thus **indicating** a *Helicobacter* caused gastrointestinal disorder.

An acidic fluid is further administered to the subject prior to administering the capsule, thus eliminating false positive readings (claimed).

Dwg.0/1

FS CPI

FA AB; DCN

MC CPI: B04-C01; B04-F10; B04-L05; B04-N03; B11-A02; B11-C08E1; B11-C08E3; B11-C09; B12-K04A4; B12-K04E; D05-A02C; D05-H04; D05-H08; D05-H09; D05-H10

TECH UPTX: 20010625

TECHNOLOGY FOCUS - BIOTECHNOLOGY - Preferred Method: Both the first



**indicator** and the second **indicator** combination are carried by a pharmaceutically acceptable dense carrier having a density greater than body fluids, the pharmaceutically acceptable dense carrier delivering the first **indicator** and the second **indicator** combination to the gastric mucosa. The dense carrier is dissolved in the gastric fluids after the soluble container is dissolved. The pharmaceutically acceptable carrier is sugar beads, and carrier has a diameter from about 0.2-3.0 mm, thus facilitating dispersal of the **indicators** over the gastric mucosa.

A first portion of the carrier is coated with the first **indicator** and a second portion of the carrier is coated with the second **indicator** combination. The first **indicator** is sorbed by a first portion of the carrier and the second **indicator** combination is sorbed by a second portion of the carrier. A buffer is added to the dense carrier in order to neutralize the **pH** of the dense carrier. The reagent is **urea**, and the **urea** reacts with the **urease** produced by *Helicobacter* to generate **ammonia**. The first **pH indicator** and the second **pH indicators** are weak acids that exhibit a first color that **indicates** an acid **pH** range and a second color that **indicates** an alkaline range.

The first **pH indicator** is bromothymol blue (dibromothymolsulfonphthalein) and the second **pH indicator** is **phenol red** (

**phenolsulfonphthalein**). (M) preferably involves:

- (a) providing at least two separate groups of pharmaceutically acceptable **pH indicator** sorbing dense carriers having a density greater than body fluids to cause the carriers to descend through the patient's gastric fluids to the patient's gastric mucosa;
- (b) combining a first of the two separate groups of dense carriers with a pharmaceutically acceptable first **pH indicator** that exhibits a first indicium when exposed to an acidic **pH** range and a second indicium when exposed to an alkaline **pH** range;
- (c) combining a second of the two separate groups of dense carriers with a combination of a pharmaceutically acceptable second **pH indicator** and **urea**, the second **pH indicator** exhibiting a first indicium when exposed to an acidic **pH** range and a third indicium when exposed to an alkaline **pH** range, the first **pH indicator** first indicium and the second **pH second indicator** first indicium being the same, the first **pH indicator** second indicium and the second **pH indicator** combination third indicium being different from one another and from the first **pH indicator** first indicium and the second **pH indicator** first indicium;
- (d) administering the first dense carrier and the second dense carrier to a patient;
- (e) contacting the patient's gastric mucosa with the first **indicator**, the second **indicator** and the **urea** contained within the carriers;
- (f) raising **pH** levels proximate to the second **pH indicator** and **urea** in response to the increased **ammonia** generated by a reaction between the **urea** and the **urease**;
- (g) observing the **indication** of **urease** producing *Helicobacter* in the patient's stomach by observing the first **pH indicator** and the second **pH indicator** combination, where:
  - (i) both the first indicium of the first **indicator** and the first indica of the second **indicator** combination **indicating** an acidic **pH** range **indicating** an absence of *Helicobacter* and that the stomach is acidic;
  - (ii) both the second indicium of the first **indicator** and the

second indicium of the second **indicator** combination **indicating** a false positive result and that the stomach is alkaline; or

(iii) the second indicium of the first **indicator** **indicating** an acidic pH range and the second indicium of the second **indicator** combination **indicating** an alkaline pH range, signifies the presence of **urease** producing *Helicobacter* and that the stomach is acidic;

(h) determining, based on observation (i), that the stomach is acidic and that there is an absence of **urease** producing *Helicobacter*, observation (ii), that the stomach is alkaline and no determination can be made, or observation (iii), that there is a presence of **urease** producing *Helicobacter* in the patient's stomach.

ABEX UPTX: 20010625

EXAMPLE - No relevant example is given.

L113 ANSWER 6 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 2000-560698 [52] WPIX

DNN N2000-415093 DNC C2000-167388

TI Measurement of **urea** nitrogen for diagnosing renal diseases, comprises detecting an optical change in pH in the liquid phase by **ammonia** formed by reacting **urea** in liquid phase containing specific buffer solutions.

DC B04 D16 S03

PA (IATR) IATRON LAB INC

CYC 1

PI JP 2000189196 A 20000711 (200052)\* 5p C12Q001-58 <--

ADT JP 2000189196 A JP 1998-376480 19981225

PRAI JP 1998-376480 19981225

IC ICM C12Q001-58

ICS G01N033-62

AB JP2000189196 A UPAB: 20001018

NOVELTY - Measurement of **urea** nitrogen comprises reacting **urea** and **urease** in liquid phase containing two or more kinds of buffer solutions and detecting optically the change in pH in liquid phase by **ammonia** formed in the reaction.

DETAILED DESCRIPTION - An INDEPENDENT CLAIM is also included for the reagent for **urea** nitrogen measurement.

USE - For diagnosing renal diseases.

ADVANTAGE - Wide range of concentration of **urea** nitrogen is determined. The liquid reagent is inexpensive and stable for long duration. The effect of measurement of **urea** nitrogen is improved.

Dwg.1/2

FS CPI EPI

FA AB; GI; DCN

MC CPI: B04-B04B1; B04-L05; B10-A01; B10-A13C; B11-C07B1;

B11-C08E3; B12-K04A; D05-A02C; D05-H09

EPI: S03-E14H

L113 ANSWER 7 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 2000-221147 [19] WPIX

DNN N2000-165428

TI Fabrication of an encapsulated pharmaceutical detecting **urease** in the stomach - for identification of *helicobacter pilori* infection by means of phenol sulphonethalein and thymol sulphone thalein reagents, with buffer and saccharose and urea modified by carbon 14. NoAbstract.

DC S03

IN MARSHALL, B J

PA (MARS-I) MARSHALL B J

CYC 1

PI MX 9703147 A1 19981001 (200019)\*

G01N033-573 <--

ADT MX 9703147 A1 MX 1997-3147 19970429

PRAI MX 1997-3147 19970429  
 IC ICM G01N033-573  
 FS EPI  
 FA NOAB  
 MC EPI: S03-E14H4

L113 ANSWER 8 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 1998-437884 [38] WPIX

DNC C1998-133216

TI Determination of *Helicobacter pylori* infection levels - by quantitative determination of ammonia production in a urea solution containing a tissue sample.

DC B04 D16

PA (DEJA-I) DEJACO R

CYC 1

PI AT 9400611 A 19980715 (199838)\* 9p C12Q001-58 <--

AT 404840 B 19990115 (199908) C12Q001-58 <--

ADT AT 9400611 A AT 1994-611 19940323; AT 404840 B AT 1994-611 19940323

FDT AT 404840 B Previous Publ. AT 9400611

PRAI AT 1994-611 19940323

IC ICM C12Q001-58

AB AT 9400611 A UPAB: 19980923

Use of a device for quantitative determination of ammonia in aqueous samples, particularly test strips with a concentration dependent colour change, for determination of levels of *Helicobacter pylori* by reaction of a tissue sample with a urea solution, such that the concentration of ammonia corresponds to that of human blood.

USE - The technique is useful for determination of levels of *H. pylori* infection.

Dwg.0/0

FS CPI

FA AB

MC CPI: B04-F10; B05-C01; B10-A13C; B11-C07B1;  
 B12-K04A4; D05-H04

L113 ANSWER 9 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 1995-283092 [37] WPIX

DNN N1995-215475 DNC C1995-127739

TI Test compsns. for detection of *Helicobacter pylori* urease - contg. urea and a combination of pH indicator dyes.

DC B04 D16 S03

IN JACKSON, F W

PA (CHEK-N) CHEK-MED SYSTEMS INC

CYC 20

PI US 5439801 A 19950808 (199537)\* 7p C12Q001-58 <--

WO 9521937 A1 19950817 (199538) EN 27p C12Q001-58 <--

RW: AT BE CH DE DK ES FR GB GR IE IT LU MC NL PT SE

W: CA JP MX

CA 2160916 C 19990330 (199931) C12Q001-58 <--

MX 9504461 A1 19990501 (200056) C12Q001-58 <--

MX 196023 B 20000414 (200124) G01N033-048 <--

ADT US 5439801 A US 1994-195954 19940214; WO 9521937 A1 WO 1995-US1608 19950206; CA 2160916 C CA 1995-2160916 19950206; MX 9504461 A1 MX 1995-4461 19951020; MX 196023 B MX 1995-4461 19950206

PRAI US 1994-195954 19940214

REP US 4748113; US 5258178; US 5260057; US 5314804

IC ICM C12Q001-58; G01N033-048

ICS A01N001-02; C12Q001-00; C12Q001-04; C12Q001-62;  
 G01N033-48

AB US 5439801 A UPAB: 19950921

Test compsns. for diagnosis of gastric disease by detection of

**urease** associated with *Helicobacter pylori* in a **biopsy** specimen contain **urea** and 2 **pH indicator** dyes such that the colour change **indicating** the presence of *H. pylori* initially occurs at an acid **pH** and the resulting colour is distinct from the colour of the **biopsy** specimen. Also claimed is a compsn. as above contg. 0.5-2 wt.% **urea**, 0.4-1.4 wt.% agar, 0.2-1.2 wt.% N-octyl glucose and 1.5-3.5 mM NaH<sub>2</sub>PO<sub>4</sub>, the balance comprising a preservative, the **indicator** dyes and water, where the compsn. is in the form of a gel soft enough to envelop a **biopsy** specimen pushed into it and has an initial acid **pH**

USE - The compsns. are esp. useful for diagnosis of peptic ulcers.

ADVANTAGE - Compared with prods. based on **phenol**

**red**, e.g. 'Clotest' (RTM), the compsns. give a more distinctive colour change at a lower **pH**, which excludes false positive due to other bacteria, e.g. *Proteus* and *Pseudomonas* spp..

Dwg.0/0

FS CPI EPI

FA AB; DCN

MC CPI: B04-C02D; B04-L05; B05-B02A3; B06-C; B10-A07; **B10-A13C**;  
B10-A16; B10-E02; **B11-C07B1**; **B12-K04A**; D05-A02C;  
D05-H04

EPI: **S03-E14H9**

L113 ANSWER 10 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN **1995-206245** [27] WPIX

CR 1993-320766 [40]

DNN **N1995-161623** DNC **C1995-095613**

TI Device for detecting *Helicobacter pylori* by measuring **urease** levels - comprises a **urease** substrate, an **ammonia**-sensitive **indicator** and sulphamic acid.

DC B04 D16 S03

IN BOGUSLASKI, R C; CARRICO, R J

PA (SERI-N) SERIM RES CORP

CYC 1

PI US 5420016 A 19950530 (199527)\* 8p C12Q001-58 <--

ADT US 5420016 A CIP of US 1992-856992 19920324, US 1994-198236 19940218

FDT US 5420016 A CIP of US 5314804

PRAI US 1994-198236 19940218; US 1992-856992 19920324

IC ICM **C12Q001-58**

ICS **C12Q001-04**; **G01N021-00**

AB US 5420016 A UPAB: 19950712

Multilayer test device for detecting **urease** in biological **tissue** specimens comprises: (a) a substrate element comprising a matrix contg. a **urease** substrate; (b) a diffusion element comprising a **NH3**-permeable and water-impermeable membrane; (c) an **indicator** element comprising a matrix contg. a **NH3**-sensitive **indicator**. The **indicator** and diffusion elements are contiguous and one contains sufficient sulphamic acid to react with **NH3** to produce a desired sensitivity. The device is designed so that the **tissue** specimen can be placed between the substrate and diffusion elements. Also claimed is a test kit comprising an aq. rehydrating soln., a buffer with a **pH** of 7-9 and a device as above where the **indicator** is the dried residue of a **pH indicator** with a pKa of 2-6.

USE - The device may be used to detect **urease**-producing microorganisms, esp. *Helicobacter pylori*, in human gastric mucosa **biopsy** specimens.

ADVANTAGE - The sulphamic acid is included to scavenge pre-existing **NH3** so that only **urease**-generated **NH3** is detected.

Dwg.1/5

FS CPI EPI

FA AB; GI; DCN  
 MC CPI: B04-B04L; B04-F10A; B04-L05; B05-C03; B11-C08E1; **B12-K04A**;  
 D05-H04  
 EPI: S03-E09E; **S03-E14H6**

L113 ANSWER 11 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN **1995-178631** [23] WPIX

CR 2001-335013 [33]

DNN **N1995-140283** DNC **C1995-082674**

TI In vivo detection of **urease**-producing *Helicobacter* - using two reagents which react differently, through colour change, to the increase in **pH**.

DC B04 D16 S03

IN **MARSHALL, B; MARSHALL, B J**

PA (MARS-I) MARSHALL B; (MARS-I) MARSHALL B J

CYC 60

PI WO 9511672 A1 19950504 (199523)\* EN 16p A61K009-28

RW: AT BE CH DE DK ES FR GB GR IE IT KE LU MC MW NL OA PT SD SE SZ

W: AM AT AU BB BG BR BY CA CH CN CZ DE DK EE ES FI GB GE HU JP KE KG

KP KR KZ LK LR LT LU LV MD MG MN MW NL NO NZ PL PT RO RU SD SE SI

SK TJ TT UA US UZ VN

AU 9481270 A 19950522 (199534) A61K009-28

EP 725633 A1 19960814 (199637) EN A61K009-28

R: AT CH DE GB IE LI LU

JP 09506246 W 19970624 (199735) 17p C12Q001-58 <--

BR 9407718 A 19971111 (199801) A61K009-28

CN 1139381 A 19970101 (199809) A61K009-28

ADT WO 9511672 A1 WO 1994-US12332 19941025; AU 9481270 A AU 1994-81270 19941025; EP 725633 A1 WO 1994-US12332 19941025; EP 1995-900448 19941025; JP 09506246 W WO 1994-US12332 19941025; JP 1995-512826 19941025; BR 9407718 A BR 1994-7718 19941025; WO 1994-US12332 19941025; CN 1139381 A CN 1994-194624 19941025

FDT AU 9481270 A Based on WO 9511672; EP 725633 A1 Based on WO 9511672; JP 09506246 W Based on WO 9511672; BR 9407718 A Based on WO 9511672

PRAI US 1993-142600 19931028

REP US 5262156; US 5314804

IC ICM A61K009-28; **C12Q001-58**

ICS A61K009-48; A61K009-54; **C12Q001-04**; **G01N021-77**

AB WO 9511672 A UPAB: 20010625

In vivo detection of **urease**-producing *Helicobacter* (I) in the upper stomach comprises: (i) obtaining at least 2 separate gps. of dense carriers; (ii) combining the first gp. with a first reagent **indicator** (R1); (iii) combining the second gp. with a combination of a second reagent **indicator** (R2) and **urea**; (iv) encapsulating R1 and the R2-**urea** combination in a soluble capsule; (v) administering the capsule to a patient; (vi) causing the capsule to migrate to the gastric mucosa through the density of the carriers; (vii) dissolving the capsule contg. R1 and R2-**urea** in the gastric juices, such that R1 and R2-**urea** are placed in direct contact with the gastric mucosa, allowing the **urea** to react with any **urease** present in the stomach, thus creating **ammonia**, the **ammonia** causing the **pH** within the stomach to increase, this causing R1 and R2 to react to the increase in **pH**, the reaction being viewed through endoscopy. A diagnostic device is also provided.

USE - The method is useful for in vivo diagnosis of upper gastrointestinal diseases, esp those mediated by infection of gastric mucosa by *Helicobacter pylori*.

ADVANTAGE - The novel method of detecting alkaline **pH** change in vivo cuts down the number of **biopsies** required and is safe for patients having any bleeding tendencies. It is also a rapid, low cost test. Additionally, through the colour change, it can be determined if the change is a true positive or a false positive reaction.

Dwg.0/1  
 FS CPI EPI  
 FA AB; GI; DCN  
 MC CPI: B04-F10; B04-L05; **B10-A13C**; **B11-C07B1**;  
**B12-K04A**; D05-H04  
 EPI: S03-E04E; **S03-E14H9**

L113 ANSWER 12 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 1993-320766 [40] WPIX

CR 1995-206245 [27]

DNN **N1993-247028** DNC **C1993-142812**

TI Detection of **urease** in human biological **tissue** - by contact with buffered **urea** and using formed **ammonia** to change colour of **indicator**, used esp. for diagnosing *Helicobacter pylori* infection.

DC B04 D16 S03

IN BOGUSLASKI, R C; CARRICO, R J

PA (SERI-N) SERIM RES CORP

CYC 20

PI WO 9319200 A1 19930930 (199340)\* 26p C12Q001-58 <--

RW: AT BE CH DE DK ES FR GB GR IE IT LU MC NL PT SE

W: AU CA JP

AU 9337361 A 19931021 (199407) C12Q001-58 <--

US 5314804 A 19940524 (199420) 7p C12Q001-58 <--

EP 633946 A1 19950118 (199507) EN

R: DE DK FR GB IT SE

JP 07505279 W 19950615 (199532) C12Q001-58 <--

EP 633946 A4 19960626 (199644) C12Q001-58 <--

JP 2638682 B2 19970806 (199736) 8p C12Q001-58 <--

CA 2131317 C 19980224 (199817) C12Q001-58 <--

EP 633946 B1 20010801 (200144) EN C12Q001-58 <--

R: DE DK FR GB IT SE

DE 69330515 E 20010906 (200159) C12Q001-58 <--

ADT WO 9319200 A1 WO 1993-US1819.19930303; AU 9337361 A AU 1993-37361

19930303; US 5314804 A US 1992-856992 19920324; EP 633946 A1 EP

1993-906267 19930303, WO 1993-US1819 19930303; JP 07505279 W JP

1993-516569 19930303, WO 1993-US1819 19930303; EP 633946 A4 EP 1993-906267

; JP 2638682 B2 JP 1993-516569 19930303, WO 1993-US1819 19930303; CA

2131317 C CA 1993-2131317 19930303; EP 633946 B1 EP 1993-906267 19930303;

WO 1993-US1819 19930303; DE 69330515 E DE 1993-630515 19930303, EP

1993-906267 19930303, WO 1993-US1819 19930303

FDT AU 9337361 A Based on WO 9319200; EP 633946 A1 Based on WO 9319200; JP

07505279 W Based on WO 9319200; JP 2638682 B2 Previous Publ. JP 07505279,

Based on WO 9319200; EP 633946 B1 Based on WO 9319200; DE 69330515 E Based

on EP 633946, Based on WO 9319200

PRAI US 1992-856992 19920324

REP EP 458231; US 3876502; US 4748113; US 4830010; US 4923801; 2.Jnl.Ref

IC ICM **C12Q001-58**

ICS C12M001-40; C12Q001-26; C12Q001-62; **G01N021-77**

ICA C12Q001-00; **C12Q001-04**

ICI **C12Q001-04**, C12R001:

AB WO 9319200 A UPAB: 20011012

**Urease** (I) is detected in a biological **tissue sample** by (i) placing the **sample** on a diffusion element permeable to **NH3**; (2) treating the **sample** with a **pH-optimised soln. of urea** substrate (II) so as to produce **NH3**; (3) allowing **NH3** to diffuse through the diffusion element so that it contacts an **indicator** element on the other side; and (4) observing reaction of **NH3** with the **indicator**.

Also new are multilayer test devices and kits for this process.

Preg. (II) is present in a substrate element, comprising a matrix and **pH 7-9 buffer**. The indica element comprises a matrix contg. a

pH-sensitive dye of pKa less than 8 (esp. 2-6).

The **sample** is placed on the diffusion element, then the substrate placed on top, partic. by folding over the support to which both elements are attached. The **indicator** or diffusion element may contain a known amt. of cpd. (esp. sulphamic acid) which reacts with any **NH3** already present (to ensure that only (I) - generated **NH3** is detected.) II Partic. the substrate element comprises absorbent paper impregnated with a buffered **urea** soln. then dried, and the diffusion element is a membrane of pore size 0.05-10 (esp. 0.1-10) microns, e.g. of PTFE. II To provide a positive control, a known amt. of **urease** is placed on the diffusion element, away from the test **sample**. USE/ADVANTAGE - The method is used to detect helicobacter pyloric (a possible cause of gastritis and ulcers) in human gastric mucosal **biopsies**. In theis test components of the **sample** do not interfere with the **indicator** reaction, and the **sample** is incubated at pH optimal for (I) activity.

Dwg. 1/4

Dwg. 1/4

FS CPI EPI

FA AB; GI; DCN

MC CPI: B04-B02C3; B10-A13D; B11-C08E3; **B12-K04A**; D05-A02C; D05-H09

EPI: S03-E04E; **S03-E14H6**

ABEQ US 5314804 A UPAB: 19940705

Detecting **urease** in a biological **tissue** specimen comprises (A) positioning the specimen on 1 side of a diffusion element permeable to **ammonia**; (B) contacting the specimen with a pH optimised **urease** substrate comprising a soln. of **urea** and a buffer having a pH of 7.0-9.0; (c) permeating the obtd. **ammonia** through the diffusion element to contact an **indicator** element at the opposite side of and contiguous with the diffusion element; and (D) observing the reaction of **ammonia** with the **indicator** element. The **indicator** element comprises a matrix contg. a pH **indicator** having a pKa of 2.0-6.0.

Pref. the diffusion element is a membrane having a pore size of 0.05-10 microns. The soln. of **urea** and buffer is contained in a matrix to form the **urease** substrate.

USE/ADVANTAGE - Used for determining the presence of Helicobacter pylori. The method is rapid and easy.

Dwg.1/4

L113 ANSWER 13 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 1991-232336 [32] WPIX

DNN N1991-177148 DNC C1991-101006

TI Measurement of **urea** or **urease** in biological fluids - by mixing with pH **indicator** and **urease** or **urea**.

DC B04 D16 J04 S03 S05

IN ORSONNEAU, J L

PA (HOSP-N) CENT HOSPIT REG UNI

CYC 1

PI FR 2654436 A 19910517 (199132)\*

ADT FR 2654436 A FR 1989-14907 19891114

PRAI FR 1989-14907 19891114

IC C12Q001-58; G01N021-79; G01N033-62

AB FR 2654436 A UPAB: 19930928

**Urea** or **urease** is measured in liqs., partic. biological fluids, by the following methods: the fluid is mixed with a first reagent contg. a stable dye the colour of which varies with pH in the range 5.5-9, it is then mixed with a second reagent contg. **urea** or **urease** which ever one is not present in the test soln.. The optical density of the mixt. is then measured at the

same wavelength of visible light before and after hydrolysis due to the action of the **urease**. The difference is compared with the result obtained with standard solns. and so the concn. of **urea** or **urease** is calculated.

ADVANTAGE - This process is cheap and simple to carry out, may be effected on urine **samples** without interference from **ammonia** present, and it does not require pre-treatment of the **sample** soln..

0/0

FS CPI EPI

FA AB; DCN

MC CPI: B04-B02C3; B04-B04B; B06-A02; **B10-A13C**; B11-C07B2;

**B12-K04A**; D05-A02C; D05-H09; J04-B01

EPI: S03-E04E; **S03-E14H**; S05-C09

L113 ANSWER 14 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN **1990-376291** [51] WPIX

DNC **C1990-163955**

TI Detection of **urease** in endoscopic **biopsies** - by colour change of **urea** soln. contg. **phenol red** indicator.

DC B04 D16 J04

IN ISERHARD, R

PA (ISER-I) ISERHARD R

CYC 1

PI BR 8902699 A 19901120 (199051)\*

ADT BR 8902699 A BR 1989-2699 19890519

PRAI BR 1989-2699 19890519

IC **C12Q001-58**

AB BR 8902699 A UPAB: 19930928

The enzyme **urease** performed in endoscopic **biopsies** of gastro-duodenal mucous membrane by bacterial action, is detected by immersing the **biopsy** specimen in a gelatinous soln. contg. peptone 1.0 g/l., glucose 1.0, sodium chloride 5, monobasic K phosphate 2, **Phenol Red** 0.012, **urea** 20, Metronidazol 0.002, Gentamicine 0.24 and agar-agar 12 g/l., in dist. water, in presence of **urease**, **ammonia** and bicarbonate are liberated, raising the pH from 5.8 to over 6.0 and changing the colour of the gel from pale yellow to red. The anti-bacterial agents prevents contamination by bacteria from **biopsy** equipment.

FS CPI

FA AB

MC CPI: B02-G; B04-B02C3; B06-C; B07-D09; **B10-A13C**;

**B11-C07B1**; **B12-K04A**; D05-H09; J04-B01

L113 ANSWER 15 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN **1990-178354** [23] WPIX

DNC **C1990-077456**

TI Enrichment and isolation of campylobacter pylori - using acidic medium to kill non-**urease** producing bacteria and plating on agar contg. selective antibiotics.

DC B04 D16 J04

IN GUERRANT, R L; **MARSHALL, B J**

PA (UYVI-N) UNIV VIRGINIA

CYC 1

PI US 4923801 A 19900508 (199023)\*

ADT US 4923801 A US 1987-37938 19870413

PRAI US 1987-37938 19870413

IC **C12Q001-58**

AB US 4923801 A UPAB: 19930928

Enrichment and isolation of campylobacter pylori from a specimen contaminated with a plurality of non-**urease** and **urease** producing organisms comprises: (a) homogenizing a specimen contaminated



with organisms in water; (b) introducing the homogenate into an acidified (pH<2.5) soln. of urea, so that most of the non-urease producing and some of the urease-producing organisms are killed by the acid medium, those remaining being pretreated from acid attack by creating a protective ammonium layer by breaking down the urea; (c) plating the remaining urease-producing organisms onto a medium contg. antibiotics inhibitory to most of these organisms but not to C. pylon; and (d) detecting the presence of colonies of C.pylon.

USE/ADVANTAGE - C.pylon is a slow growing fastidious organisms and could not previously be easily isolated from biological specimens contg. contaminating bacteria. The new method of isolation utilizes the discovery that C pylon is able to survive in acid medium provided that urea is present, by prodn. of urease which breaks down the urea to ammonia which neutralises the acid and protects th organism. It is therefore possible to isolate C. pylon from leading contaminated specimens such as stool. Early detection and isolation of C. pylon would enable specific etiological diagnosis of this infection, and rapid determination of antibiotic sensitivities. @  
0/0

FS CPI

FA AB; DCN

MC CPI; B04-B02B1; B11-C08E3; **B12-K04A4**; D05-A02C; D05-H06; J04-B01

L113 ANSWER 16 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 1986-340935 [52] WPIX

CR 1986-340936 [52]

DNC C1986-147787

TI Treatment of gastrointestinal disorders - with daily dosages of bismuth e.g. in salt form.

DC B05 B06 P32

IN **MARSHALL, B J**

PA (MARS-I) MARSHALL B J; (PROC) PROCTER &amp; GAMBLE CO

CYC 14

PI EP 206626 A 19861230 (198652)\* EN 11p

R: AT BE CH DE FR GB IT LI LU NL SE

BE 904922 A 19861215 (198701)

DE 3619733 A 19870212 (198707)

DE 3619734 A 19870212 (198707)

JP 62048624 A 19870303 (198714)

EP 206626 B1 19920812 (199233) EN 6p A61K033-00

R: AT BE CH DE FR GB IT LI LU NL SE

DE 3686361 G 19920917 (199239)

JP 07094391 B2 19951011 (199545) 5p A61K031-29

PH 26891 A 19921103 (199635) A61K045-00

US 5601848 A 19970211 (199712) 6p A61K033-24

EP 206626 B2 20020522 (200241) EN A61K033-24

R: AT BE CH DE FR GB IT LI LU NL SE

ADT EP 206626 A EP 1986-304408 19860610; BE 904922 A BE 1986-904922 19860613;

DE 3619733 A DE 1986-3619733 19860612; DE 3619734 A DE 1986-3619734

19860612; JP 62048624 A JP 1986-138038 19860613; EP 206626 B1 EP

1986-304408 19860610; DE 3686361 G DE 1986-3686361 19860610, EP

1986-304408 19860610; JP 07094391 B2 JP 1986-138038 19860613; PH 26891 A

PH 1986-33888 19860713; US 5601848 A Cont of US 1985-744842 19850613, US

1987-70857 19870708; EP 206626 B2 EP 1986-304408 19860610

FDT DE 3686361 G Based on EP 206626; JP 07094391 B2 Based on JP 62048624

PRAI US 1985-744842 19850613; US 1987-70857 19870708

REP 3.Jnl.Ref; A3...8915; EP 75992; FR 5877; FR 6197; GB 1107655; No-SR.Pub;  
US 3577533; 8.Jnl.RefIC A61D000-00; A61K031-19; A61K031-29; A61K033-24; **C12Q001-58**;  
G01N033-50

ICM A61K031-29; A61K033-00; A61K033-24; A61K045-00

ICS A61D000-00; A61K031-19; A61K031-60; **C12Q001-58**; G01N033-50

AB EP 206626 A UPAB: 20020701

Bismuth is used for the mfr. of a medicament for treatment of gastrointestinal disorders in humans and animals, the treatment comprising administration of 50,5,000 mg/day of bismuth, for 3-21 days. Pref. the daily dose of bismuth is 500-1,500 mg. The bismuth is pref. in the form of a salt e.g. the aluminate, sub-carbonate, (sub)citrate, sub-galate, sub-nitrate, tartrate or sub-salicylate or tripotassium dicitrato bismuthate. The stated wt. quantities are of elemental bismuth, so the actual wt. of bismuth contg. cpd. will be greater. Pref. before treatment with the bismuth the subject is positively diagnosed for CLO infection. Diagnosis is then repeated during the administration and administration is terminated when diagnosis yields a negative result. compsns. for administering the bismuth are conventional.

ADVANTAGE - The method cures, or affords lower relapse rates of, gastritis and peptic ulcer disease. It does not render the patient hypochlorhydric.

0/0

Dwg.0/0

FS CPI GMPI

FA AB

MC CPI: B05-A02; B12-E08; B12-J01

ABEQ DE 3686361 G UPAB: 19930922

Bismuth is used for the mfr. of a medicament for treatment of gastrointestinal disorders in humans and animals, the treatment comprising administration of 50,5,000 mg/day of bismuth, for 3-21 days. Pref. the daily dose of bismuth is 500-1,500 mg. The bismuth is pref. in the form of a salt e.g. the aluminate, sub-carbonate, (sub)citrate, sub-galate, sub-nitrate, tartrate or sub-salicylate or tripotassium dicitrato bismuthate. The stated wt. quantities are of elemental bismuth, so the actual wt. of bismuth contg. cpd. will be greater. Pref. before treatment with the bismuth the subject is positively diagnosed for CLO infection. Diagnosis is then repeated during the administration and administration is terminated when diagnosis yields a negative result. compsns. for administering the bismuth are conventional.

ADVANTAGE - The method cures, or affords lower relapse rates of, gastritis and peptic ulcer disease. It does not render the patient hypochlorhydric.

ABEQ DE 3686362 G UPAB: 19930922

Bismuth is used for the mfr. of a medicament for treatment of gastrointestinal disorders in humans and animals, the treatment comprising administration of 50,5,000 mg/day of bismuth, for 3-21 days. Pref. the daily dose of bismuth is 500-1,500 mg. The bismuth is pref. in the form of a salt e.g. the aluminate, sub-carbonate, (sub)citrate, sub-galate, sub-nitrate, tartrate or sub-salicylate or tripotassium dicitrato bismuthate. The stated wt. quantities are of elemental bismuth, so the actual wt. of bismuth contg. cpd. will be greater. Pref. before treatment with the bismuth the subject is positively diagnosed for CLO infection. Diagnosis is then repeated during the administration and administration is terminated when diagnosis yields a negative result. compsns. for administering the bismuth are conventional.

ADVANTAGE - The method cures, or affords lower relapse rates of, gastritis and peptic ulcer disease. It does not render the patient hypochlorhydric.

ABEQ EP 206626 B UPAB: 19930922

The use of bismuth for the manufacture of a medicament for the treatment of a disorder of the upper gastrointestinal tract of a human or other animal subject in which the disorder is caused or mediated by *Campylobacter pyloridis*, and wherein is excluded the use of bismuth in the form of bismuth subsalicylate.

0/0

ABEQ EP 206627 B UPAB: 19930922

The use of bismuth subsalicylate for the manufacture of a medicament for the treatment of a disorder of the upper gastrointestinal tract of a human

or other animal subject in which the disorder is caused or mediated by *Campylobacter pyloridis*.

0/0

ABEQ US 5601848 A UPAB: 19970320

Treatment of a human or lower animal subject having an infectious gastrointestinal disorder caused or mediated by *Campylobacter pyloridis* comprises combating said *Campylobacter pyloridis* infection in said subject, comprising the step of orally administering to said subject from about 50 mg to about 5000 mg of bismuth, per day, for from 3 to 56 days, wherein said bismuth is selected from the group consisting of bismuth aluminate, bismuth subcarbonate, bismuth citrate, bismuth subgalate, bismuth subnitrate, bismuth tartrate, bismuth subsalicylate, and mixtures thereof.

Dwg.0/0

L113 ANSWER 17 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 1986-327235 [50] WPIX

DNN N1986-244169 DNC C1986-141647

TI Compsn. for diagnosis of gastrointestinal disorders - e.g. mediated by *Campylobacter pyloridis* infection, comprises **urea**, bactericide, **pH indicator** and water.

DC B04 D16 S03

IN MARSHALL, B J

PA (MARS-I) MARSHALL B J

CYC 20

PI EP 204438 A 19861210 (198650)\* EN 25p

R: AT BE CH DE FR GB IT LI LU NL SE

AU 8657398 A 19861120 (198702)

NO 8601966 A 19861215 (198705)

DK 8602283 A 19861118 (198707)

BR 8602243 A 19870113 (198708)

JP 62026000 A 19870203 (198710)

ZA 8603605 A 19871116 (198808)

US 4748113 A 19880531 (198824)

CA 1274757 A 19901002 (199045)

EP 204438 B 19910306 (199110)

R: AT BE CH DE FR GB IT LI LU NL SE

DE 3677820 G 19910411 (199116)

JP 06095960 B2 19941130 (199501)

6p C12Q001-58 <--

KR 9406322 B1 19940716 (199617)#

A61K031-17

DK 173710 B 20010709 (200147)

C12Q001-58 <--

ADT EP 204438 A EP 1986-303493 19860508; JP 62026000 A JP 1986-112427

19860516; ZA 8603605 A ZA 1986-3605 19860515; US 4748113 A US 1985-744840

19850613; JP 06095960 B2 JP 1986-112427 19860516; KR 9406322 B1 KR

1986-7444 19860905; DK 173710 B DK 1986-2283 19860516

FDT JP 06095960 B2 Based on JP 62026000; DK 173710 B Previous Publ. DK 8602283

PRAI US 1985-744840 19850613; KR 1986-7444 19860905

REP 2.Jnl.Ref; A3...8721; EP 18825; FR 2442268; GB 1112251; JP 58077663;

No-SR.Pub; US 3145086; US 4101382; US 4282316; JP 58077172

IC C12M001-34; C12Q001-58; G01N033-00

ICM A61K031-17; C12Q001-58

ICS A61B005-00; C12M001-34; G01N033-00

ICA G01N033-62

AB EP 204438 A UPAB: 19930922

Compsn. for detection of preformed **urease** comprises **urea**, a bactericide, sufficient **pH indicator** undergoing a colour change on increase of **pH**, and water. The comps. has acid **pH** of at least 5.0 and **pH** is at least 1 **pH** unit lower than the pKa of the **indicator**.

Device for use as above comprises a container of vol. 40-1000 cu.mm. with an aperture area of 20-200 sq.mm, with a movable cover to open and close the opening, and contg. 0.04-2.0 pref. 0.2-0.4 ml of the above compsn..

USE/ADVANTAGE - Useful in diagnosis of gastrointestinal diseases mediated by e.g. *Campylobacter pyloridis*, which produces a high activity **urease**. Compsn. gives rapid, inexpensive and accurate diagnosis, of e.g. chronic or atrophic gastritis, gastroenteritis, dyspepsia, oesophageal reflux disease, gastric and duodenal ulcers, etc.. The bactericide ensures that only preformed **urease** is analysed.

3/3

FS CPI EPI

FA AB

MC CPI: B04-B02C3; B06-C; **B10-A13C**; **B11-C07B1**;  
**B12-K04A**; B12-K04D; D05-A02C; D05-H09  
EPI: **S03-E14H9**

ABEQ EP 204438 B UPAB: 19930922

A composition for the diagnosis of gastrointestinal disorder in a human or lower animal subject by detection of **urease** in gastric material of the subject characterised in that it comprises (a) **urea**; (b) a bactericide which substantially inhibits growth of **urease** producing organisms; (c) a **pH indicator** which undergoes a colour change upon an increase of **pH**, at an effective concentration; and (d) water; wherein said composition has an acid **pH** of at least 5.0 and the **pH** of said composition is at least about one **pH** unit lower than the pKa of said **indicator**.

ABEQ US 4748113 A UPAB: 19930922,

Compsn. for diagnosis of gastrointestinal disorders, by detecting **urease** in gastric material of the patient, comprises (a) 10-40 g/l **urea**, (b) 1-5 g/l bactericide to inhibit growth of **urease** -producing organisms, (c) **indicator** having pKa 6.5-8.5, and (d) water. The comps. has **pH** 5.0-6.5 and the **pH** is at least 1 unit below the pKa of the **indicator**.

The **indicator** is e.g. **phenol red**. The comps. opt. includes a buffer and a gelling agent, e.g. non-nutritive agar.

ADVANTAGE - Compsn. allows rapid, inexpensive and accurate diagnosis of disorders of the upper gastrointestinal tract.

L113 ANSWER 18 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN 1986-091422 [14] WPIX

DNN N1986-066659 DNC C1986-039132

TI Colorimetric determ. of **ammonia** concn. - formed by **urease** treatment of **urea**, using phenol deriv. and oxidising agent in presence of imidazole or its deriv..

DC B04 D16 J04 S03

PA (WAKP) WAKO PURE CHEM IND LTD

CYC 1

PI JP 61038463 A 19860224 (198614)\* 11p

PRAI JP 1984-160143 19840730

IC G01N033-50

AB JP 61038463 A UPAB: 19930922

The process involves colorimetry of **ammonia** using phenol system cpd. and oxidising agent. Colorimetry is carried out in presence of imidazole and/or an imidazole deriv.

Imidazole deriv. is e.g. 1-methylimidazole, 1-ethylimidazole, 1-phenylimidazole, 1-benzylimidazole, 2-methylimidazole, 2-ethylimidazole, 2-phenylimidazole, 1,2-dimethylimidazole, etc. The concn. of imidazole and/or imidazole deriv. in the final coloured liq. is more than 1 m mol./l., pref. 1-100 m mol./l. Phenol cpd. used as colouring component in the method is e.g. phenol, salicylic acid, m-hydroxybenzoic acid, p-hydroxybenzoic acid, sulphosalicylic acid, O-, m- or p-cresol, o-methoxyphenol, etc. N,N-disubstd. aniline cpd. such as N,N-dimethylaniline, N,N-dimethyl-m-toluidine, etc. may be used as colouring component in place of phenol system cpd. The concn. of phenol system cpd. of N,N-disubstd. aniline cpd. in the final coloured liq. is

more than 10 m mol./l., pref. 50-500 m mol/l. Oxidising agent used is e.g. hypochlorite, dichloroisocyanurate, chloramine T, etc., which is used in a concn. of effective chlorine of more than 0.01%, pref. 0.03-0.3%, in the final coloured liq.

USE/ADVANTAGE - For determ. of **ammonia** formed by treating **urea** with **urease**. Imidazole or imidazole deriv. used in the method is non-toxic, not harmful and stable to light in contrast with sodium nitroprusside previously used. Consequently, prepn. of **sample** liq. is easily carried out and the stability of imidazole or its deriv. in a liq. of **urease** used for the determ. of **urea** in blood serum is very high. Also as the colouring sensitivity of the colorimetry in the presence of imidazole or its deriv. is about 1/20 times that in the previous case of using sodium nitroprusside, the determ. of blood serum **urea** with **urease** is carried out at the absorption wavelength and the colorimetry is not affected by coloured substances in blood serum, such as haemoglobin, etc.

0/0

FS CPI EPI

FA AB

MC CPI: B04-B04D4; B05-C01; B07-D09; **B10-A13C**; B10-C03; B10-E02;

B11-C08; **B12-K04**; D05-H08; J04-B01B

EPI: **S03-E14H**

L113 ANSWER 19 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN **1983-59481K** [25] WPIX

DNN **N1983-107229** DNC **C1983-057712**

TI **Urea** analysis by enzyme hydrolysis of **urea** in **sample** - converting resulting ammonium carbonate into **ammonia** and measuring amt. of **ammonia** by **indicator** discolouration.

DC B04 D16

PA (KYOT-N) KYOTO DAIICHI KAGAKU KK

CYC 1

PI JP 58077663 A 19830511 (198325)\* 8p

PRAI JP 1981-177660 19811102

IC **C12Q001-58; G01N033-62**

AB JP 58077663 A UPAB: 19930925

Method comprises (a) hydrolysing **urea** in a **sample** by an enzyme system exhibiting **urease** activity in a **sample** hole which can be kept air-tight, (b) converting resulting ammonium carbonate into **ammonia** gas under alkaline condition, (c) leading the gas via a gas-permeable membrane into an **indicator** layer, and (d) determining the concn. of **urea** based on discoloration of the **indicator** corresp. to change in pH by **ammonia** gas.

**Urea** in biological fluids such as blood, blood serum, blood plasma, saliva, etc. can be rapidly, accurately and precisely determined regardless of kind of **sample** liq.

FS CPI

FA AB

MC CPI: B04-B04B; B04-B04D; B04-B04G; **B10-A13C**; **B11-C07B**;

**B12-K04**; D05-A02

L113 ANSWER 20 OF 20 WPIX (C) 2003 THOMSON DERWENT

AN **1979-86085B** [48] WPIX

TI Diagnostic agent for **urea** determination - comprising a **urease** reaction layer and an **ammonia** **indicator** layer on a carrier acting as hand grip.

DC B04 S03 S05

IN LANGE, H R; ROTHE, A; SELLE, A K

PA (BOEF) BOEHRINGER MANNHEIM GMBH

CYC 18

PI DE 2821469 A 19791122 (197948)\*  
 EP 5519 A 19791128 (197948) DE  
 R: AT BE CH DE FR GB IT LU NL SE  
 BR 7903035 A 19791204 (197951)  
 DK 7901995 A 19791210 (198002)  
 JP 54151096 A 19791127 (198002)  
 FI 7901533 A 19800131 (198009)  
 ZA 7902362 A 19800415 (198030)  
 US 4223089 A 19800916 (198040)  
 DD 143662 A 19800903 (198046)  
 CA 1113356 A 19811201 (198201)  
 EP 5519 B 19830413 (198316) DE  
 R: AT BE CH DE FR GB IT LU NL SE  
 DE 2965181 G 19830519 (198321)  
 JP 62043500 B 19870914 (198740)  
 PRAI DE 1978-2821469 19780517  
 REP 2.Jnl.Ref; DE 1240306; DE 1245619; DE 2118455; DE 2249647; DE 2626367; DE 2748857; US 2632761  
 IC C07D213-20; C07D215-00; C12Q001-58; G01N021-06; G01N031-14; G01N033-16  
 AB DE 2821469 A UPAB: 19930901  
 New diagnostic agents for the determination of **urea** consist of a hand grip affixed to which is an **indicator** layer for gaseous **ammonia**, and a reaction layer contg. a **urease** and an alkaline buffer. The **indicator** layer is firmly bound to the hand grip, while the reaction layer is held above it at a distance of 10-200  $\mu$ m maintained by means of a spacer, the spacer and the reaction layer being readily separable from the **indicator** layer.  
 Used for determination of **urea** in body fluids in the diagnosis and monitoring of kidney disorders. The agent is simple to use without additional reaction chambers, and permits determination to be carried out on small specimens of whole blood, as well as serum or plasma. Reaction time is  $\geq 10$  mins. at room temp. Visual evaluation gives semi-quantitative results, while re-emission photometry gives quantitative determination of **urea**.  
 FS CPI EPI  
 FA AB  
 MC CPI: B04-B04D; B04-C02; B04-C03; B07-D04; B10-A13C; B11-C07B; B12-K04

=> d his

(FILE 'HOME' ENTERED AT 09:08:06 ON 30 JUN 2003)  
 SET COST OFF

FILE 'REGISTRY' ENTERED AT 09:08:22 ON 30 JUN 2003  
 E UREASE/CN

L1 1 S E3  
 L2 558 S UREASE  
 L3 557 S L2 NOT L1  
 L4 1 S UREA/CN  
 L5 1 S AMMONIA/CN  
 L6 1 S PHENOL RED/CN  
 L7 23 S 143-74-8/CRN  
 L8 7 S L7 NOT (PMS OR MXS)/CI  
 L9 5 S L8 NOT (C6-C6/ES OR CYCLODEXTRIN)  
 L10 1 S AGAR/CN

FILE 'HCAPLUS' ENTERED AT 09:11:22 ON 30 JUN 2003

L11 6909 S L1  
 L12 150 S L3  
 L13 10252 S UREASE

L14 10655 S L11-L13  
 L15 64068 S L4  
 L16 189239 S UREA  
 L17 115046 S L5  
 L18 333095 S AMMONIA OR NH3  
 L19 4892 S L14 AND L15,L16  
 L20 1587 S L19 AND L17,L18  
 L21 1287 S L6 OR L9  
 L22 2884 S PHENOL RED  
 L23 674 S PHENOLSULFONEPHTHALEIN OR PHENOLSULPHONEPHTHALEIN OR PHENOL()  
 L24 239 S PHENOLSULFONPHTHALEIN OR PHENOLSULPHONPHTHALEIN OR PHENOL() (S  
 L25 17 S L20 AND L21-L24  
 L26 5068 S L10  
 L27 47051 S AGAR  
 L28 20 S L20 AND L26,L27  
 L29 2 S L25 AND L28  
 L30 33 S L25,L28 NOT L29  
 L31 20 S L30 AND L11  
 L32 20 S L31 AND L11-L31  
 SEL DN AN 6-9 12-15 20  
 L33 11 S L32 NOT E1-E27  
 L34 13 S L30 NOT L31  
 SEL DN AN 1 5 9  
 L35 3 S L34 AND E28-E36  
 L36 16 S L29,L33,L35  
 E MCMICHAEL D/AU  
 L37 5 S E6  
 E MC MICHAEL D/AU  
 E PETERSON K/AU  
 L38 145 S E3-E17  
 E PETERSON KRIS/AU  
 L39 6 S E3,E11  
 E MARSHALL B/AU  
 L40 73 S E3,E12  
 L41 14 S E25-E27  
 E MENDIS A/AU  
 L42 23 S E3,E4,E6,E7  
 E CHAIRMAN S/AU  
 L43 1 S E4  
 E KIMBER/PA,CS  
 L44 1950 S E4-E79  
 L45 4 S E93-E100  
 L46 8 S L14 AND L37-L45  
 L47 20 S L36,L46 AND L11-L46  
 L48 10 S L33 NOT MERCURY/TI  
 L49 19 S L29,L35,L46,L48 AND L11-L48  
 L50 1122 S L1 (L) (ANST OR ANT OR DGN)/RL  
 L51 26 S L50 AND L21-L24  
 E COLOR/CT  
 E E58+ALL  
 E E2+ALL  
 L52 753 S E4,E3+NT  
 L53 2940 S E8+NT  
 L54 15989 S E2+NT  
 E E8+ALL  
 L55 803 S E6  
 L56 754 S E13+NT  
 L57 9551 S E12+NT  
 L58 44 S L50 AND L52-L57  
 L59 58 S L51,L58  
 L60 42 S L59 AND L15,L16  
 L61 10 S L59 AND L17,L18  
 L62 9 S L60 AND L61

L63 9 S L62 AND L11-L62  
L64 1 S L61 NOT L63  
L65 24 S L49,L63  
L66 32 S L60 NOT L65  
SEL DN AN 1 9 11 14 17 19 20 21 31 32  
L67 10 S E1-E30  
L68 34 S L65,L67 AND L11-L67  
L69 34 S L68 AND (UREASE OR UREA OR AMMONI? O NH3 OR NH4 OR BUFFER? OR

FILE 'HCAPLUS' ENTERED AT 09:54:14 ON 30 JUN 2003  
SEL HIT RN

FILE 'REGISTRY' ENTERED AT 09:54:32 ON 30 JUN 2003  
L70 6 S E31-E36  
L71 6 S L70 AND L1-L10

FILE 'HCAPLUS' ENTERED AT 09:55:45 ON 30 JUN 2003  
L72 1 S US20030082661/PN  
L73 1 S L72 AND L69

FILE 'WPIX' ENTERED AT 09:56:11 ON 30 JUN 2003  
L74 0 S US20030082661/PN  
L75 1083 S L13/BIX  
E UREASE/DCN  
L76 321 S L75 AND G01N/IC,ICM,ICS,ICA,ICI  
L77 242 S L75 AND G01N033/IC,ICM,ICS,ICA,ICI  
L78 30 S L75 AND G01N033-48/IC,ICM,ICS,ICA,ICI  
L79 44 S L75 AND G01N033-4?/IC,ICM,ICS,ICA,ICI  
L80 44 S L78,L79  
SEL DN AN 2 24  
L81 2 S L80 AND E1-E6  
L82 226 S C12Q001-58/IC,ICM,ICS,ICA,ICI  
L83 1200 S L75,L82  
L84 158 S L83 AND Q505/M0,M1,M2,M3,M4,M5,M6  
L85 28 S C12Q001-04/IC,ICM,ICS,ICA,ICI AND L83  
L86 178 S L84,L85  
L87 72 S L83 AND (B11-C07B OR C11-C07B OR B11-C07B1 OR C11-C07B1)/MC  
L88 203 S L86,L87  
L89 18 S L83 AND (L22/BIX OR L23/BIX OR L24/BIX)  
E PHENOL RED/DCN  
E E3+ALL  
L90 13 S L83 AND E2  
L91 207 S L88-L90  
L92 93 S L91 AND S03-E14H?/MC  
L93 162 S L91 AND (B12-K04 OR C12-K04 OR B12-K04A OR C12-K04A OR B12-K0  
L94 1 S L91 AND (B12-J01 OR C12-J01 OR B14-E10 OR C14-E10 OR B12-J OR  
L95 165 S P831/M0,M1,M2,M3,M4,M5,M6 AND L86  
L96 112 S L91 AND UREA/BIX  
E UREA/DCN  
E E3+ALL  
L97 85 S L91 AND (E2 OR 0123/DRN)  
L98 72 S L91 AND (B10-A13C OR C10-A13C)/MC  
L99 2 S L91 AND (B10-A13 OR C10-A13)/MC  
L100 122 S L96-L99  
L101 50 S L100 AND (AMMONIA OR NH3)/BIX  
E AMMONIA/DCN  
E E3+ALL  
L102 13 S L100 AND (E2 OR 1713)/DRN  
L103 50 S L101,L102  
L104 50 S L103 AND L91-L94  
SEL DN AN 1 6 7 9 12 15 16 18 22 24 25 32 38 48  
L105 14 S L104 AND E1-E35  
L106 16 S L81,L105



L107           E MCMICHAEL D/AU  
          1 S E5  
          E MC MICHAEL D/AU  
          E PETERSON K/AU  
L108           180 S E3-E22  
              E MARSHALL B/AU  
L109           41 S E3,E11  
              E MENDIS A/AU  
              E CHAIRMAN S/AU  
L110           8 S L107-L109 AND L83  
L111           20 S L106,L110 AND L75-L110  
L112           18 S L111 AND (INDICAT? OR PH OR BIOPS? OR SAMPL? OR TISSU?)/BIX  
L113           20 S L111,L112

FILE 'WPIX' ENTERED AT 10:27:58 ON 30 JUN 2003



Creation date: 12-17-2003  
Indexing Officer: ICHARLES - IRENE CHARLES  
Team: OIPEBackFileIndexing  
Dossier: 09977667

Legal Date: 08-26-2003

No.	Doccode	Number of pages
1	SRNT	14

Total number of pages: 14

Remarks:

Order of re-scan issued on .....